

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/004021 A1

(51) International Patent Classification⁷: A61K 39/395,
C07K 16/32

(21) International Application Number: PCT/US01/15677

(22) International Filing Date: 16 May 2001 (16.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/217,706 12 July 2000 (12.07.2000) US
09/772,938 31 January 2001 (31.01.2001) US

(71) Applicant: IDEC PHARMACEUTICALS CORPORATION [US/US]; 11011 Torreyana Road, San Diego, CA 92121 (US).

(72) Inventors: HANNA, Nabil; 14770 Avenida Insurgentes, Rancho Santa Fe, CA 92067 (US). HARIHARAN, Kandasamy; 7414 Park Village Road, San Diego, CA 92129 (US).

(74) Agents: TESKIN, Robin, L. et al.; Pillsbury Winthrop LLP, 1600 Tysons Boulevard, McLean, VA 22102 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(48) Date of publication of this corrected version:
16 January 2003

(15) Information about Correction:
see PCT Gazette No. 03/2003 of 16 January 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/004021 A1

(54) Title: TREATMENT OF B CELL MALIGNANCIES USING COMBINATION OF B CELL DEPLETING ANTIBODY AND IMMUNE MODULATING ANTIBODY RELATED APPLICATIONS

(57) Abstract: A combination antibody therapy for treating B cell malignancies using an immunoregulatory antibody, especially an anti-B7, anti-CD23, or anti-CD40L antibody and a B cell depleting antibody, especially anti-CD19, anti-CD20, anti-CD22 or anti-CD37 antibody is provided. Preferably, the combination therapy will comprise anti-B7 and anti-CD20 antibody administration.

Best Available Copy

**TREATMENT OF B CELL MALIGNANCIES USING
COMBINATION OF B CELL DEPLETING ANTIBODY AND IMMUNE
MODULATING ANTIBODY RELATED APPLICATIONS**

5 **Cross Reference to Related Applications**

This application claims priority from U.S. Serial Nos. 60/217,706, filed July 12, 2000, and U.S. Serial No. 09/772,938, filed January 31, 2001, and are incorporated by reference in their entirety therein.

10 **Field of the Invention**

The invention relates to a synergistic combination antibody therapy for treatment of B cell malignancies, especially B cell lymphomas and leukemias. This synergistic antibody combination comprises at least one antibody having substantial B cell depleting activity (e.g., an anti-CD19, CD20, CD22 or CD37
15 antibody) and an antibody that modulates or regulates the immune system, e.g., by modulating B cell/T cell interactions and/or B cell activity, differentiation or proliferation (e.g., anti-B7, anti-CD40, anti-CD23 or anti-CD40L). In particular, the invention encompasses combination antibody therapies for CD40+ malignancies, which include using anti-CD40L antibodies to prevent
20 CD40L from binding to CD40. These antibodies or other agents which can inhibit CD40/CD40L interaction further can be combined with chemotherapeutics, radiation and/or other antibodies, preferably B cell depleting antibodies, e.g., anti-CD19, anti-CD20, anti-CD22 and/or anti-CD40 antibodies, or fragments thereof.

25 **Background of Invention**

The immune system of vertebrates (for example, primates, which include humans, apes, monkeys, etc.) consists of a number of organs and cell types which have evolved to: accurately and specifically recognize foreign
30 microorganisms ("antigen") which invade the vertebrate-host; specifically bind to such foreign microorganisms; and, eliminate/destroy such foreign microorganisms. Lymphocytes, as well as other types of cells, are critical to the immune system and to the elimination and destruction of foreign

Best Available Copy

5 microorganisms. Lymphocytes are produced in the thymus, spleen and bone marrow (adult) and represent about 30% of the total white blood cells present in the circulatory system of humans (adult). There are two major sub-populations of lymphocytes: T cells and B cells. T cells are responsible for cell mediated immunity, while B cells are responsible for antibody production (humoral
10 immunity). However, T cells and B cells can be considered interdependent -- in a typical immune response, T cells are activated when the T cell receptor binds to fragments of an antigen that are bound to major histocompatibility complex ("MHC") glycoproteins on the surface of an antigen presenting cell; such activation causes release of biological mediators ("interleukins" or "cytokines")
15 which, in essence, stimulate B cells to differentiate and produce antibody ("immunoglobulins") against the antigen.

Each B cell within the host expresses a different antibody on its surface - thus one B cell will express antibody specific for one antigen, while another B cell will express antibody specific for a different antigen. Accordingly, B cells
20 are quite diverse, and this diversity is critical to the immune system. In humans, each B cell can produce an enormous number of antibody molecules (i.e., about 10^7 to 10^8). Such antibody production most typically ceases (or substantially decreases) when the foreign antigen has been neutralized. Occasionally, however, proliferation of a particular B cell will continue unabated; such
25 proliferation can result in a cancer referred to as "B cell lymphoma."

Non-Hodgkin's lymphoma is one type of lymphoma that is characterized by the malignant growth of B lymphocytes. According to the American Cancer Society, an estimated 54,000 new cases will be diagnosed, 65% of which will be classified as intermediate- or high-grade lymphoma.
30 Patients diagnosed with intermediate-grade lymphoma have an average survival rate of two to five years, and patients diagnosed with high-grade lymphoma survive an average of six months to two years after diagnosis.

Conventional therapies have included chemotherapy and radiation, possibly accompanied by either autologous or allogeneic bone marrow or stem
35 cell transplantation if a suitable donor is available, and if the bone marrow

5 contains too many tumor cells upon harvesting. While patients often respond to conventional therapies, they usually relapse within several months.

It is known that B cell malignancies, e.g., B cell lymphomas and leukemias may be successfully treated using antibodies specific to B cell antigens that possess B cell depleting activity. Examples of B cell antibodies
10 that have been reported to possess actual or potential application for the treatment of B cell malignancies include antibodies specific to CD20, CD19, CD22, CD37 and CD40.

Also, the use anti-CD37 antibodies having B cell depleting activity have been well reported to possess potential for treatment of B cell lymphoma. See
15 e.g., Presr et al., *J. Clin. Oncol.* 7(8): 1027-1038 (August 1989); Grossbard et al., *Blood* 8(4): 863-876 (August 15, 1992).

A. Anti-CD20 Antibodies

CD20 is a cell surface antigen expressed on more than 90% of B-cell lymphomas, which does not shed or modulate in the neoplastic cells
20 (McLaughlin et al., *J. Clin. Oncol.* 16: 2825-2833 (1998b)). The CD20 antigen is a non-glycosylated, 35 kDa B-cell membrane protein involved in intracellular signaling, B-cell differentiation and calcium channel mobilization (Clark et al., *Adv. Cancer Res.* 52: 81-149 (1989); Tedder et al., *Immunology Today* 15: 450-454 (1994)). The antigen appears as an early marker of the human B-cell
25 lineage, and is ubiquitously expressed at various antigen densities on both normal and malignant B-cell populations. However, the antigen is absent on fully, mature B-cells (e.g., plasma cells), early B-cell populations and stem cells, making it a suitable target for antibody mediated therapy.

Anti-CD20 antibodies have been prepared for use both in research and
30 therapeutics. One anti-CD20 antibody is the monoclonal B1 antibody (U.S. Patent No. 5,843,398). Anti-CD20 antibodies have also been prepared in the form of radionuclides for treating B-cell lymphoma (e.g., ¹³¹I-labeled anti-CD20 antibody), as well as a ⁸⁹Sr-labeled form for the palliation of bone pain caused by prostate and breast cancer metastasises (Endo, *Gan To Kagaku Ryoho* 26:
35 744-748 (1999)).

5 A murine monoclonal antibody, 1F5, (an anti-CD20 antibody) was reportedly administered by continuous intravenous infusion to B-cell lymphoma patients. However, extremely high levels (>2 grams) of 1F5 were reportedly required to deplete circulating tumor cells, and the results were described as "transient" (Press *et al.*, *Blood* 69: 584-591 (1987)). A potential problem with
10 using monoclonal antibodies in therapeutics is non-human monoclonal antibodies (*e.g.*, murine monoclonal antibodies) typically lack human effector functionality, *e.g.*, they are unable to, *inter alia*, mediate complement dependent lysis or lyse human target cells through antibody-dependent cellular toxicity or Fc-receptor mediated phagocytosis. Furthermore, non-human monoclonal
15 antibodies can be recognized by the human host as a foreign protein; therefore, repeated injections of such foreign antibodies can lead to the induction of immune responses leading to harmful hypersensitivity reactions. For murine-based monoclonal antibodies, this is often referred to as a Human Anti-Mouse Antibody response, or "HAMA" response. Additionally, these
20 "foreign" antibodies can be attacked by the immune system of the host such that they are, in effect, neutralized before they reach their target site.

RITUXAN®. RITUXAN® (also known as Rituximab, MabThera®, IDEC-C2B8 and C2B8) was the first FDA-approved monoclonal antibody and was developed at IDEC Pharmaceuticals (see U.S. Patent Nos. 5,843,439;
25 5,776,456 and 5,736,137) for treatment of human B-cell lymphoma (Reff *et al.*, *Blood* 83: 435-445 (1994)). RITUXAN® is a chimeric, anti-CD20 monoclonal (MAb) which is growth inhibitory and reportedly sensitizes certain lymphoma cell lines for apoptosis by chemotherapeutic agents *in vitro* (Demidem *et al.*, *Cancer Biotherapy & Radiopharmaceuticals* 12: 177- (1997)). RITUXAN®
30 also demonstrates anti-tumor activity when tested *in vivo* using murine xenograft animal models. RITUXAN® efficiently binds human complement, has strong FcR binding, and can efficiently kill human lymphocytes *in vitro* via both complement dependent (CDC) and antibody-dependent (ADCC) mechanisms (Reff *et al.*, *Blood* 83: 435-445 (1994)). In macaques, the antibody
35 selectively depletes normal B-cells from blood and lymph nodes.

5 RITUXAN® has been recommended for treatment of patients with low-grade or follicular B-cell non-Hodgkin's lymphoma (McLaughlin *et al.*, *Oncology (Huntingt)* 12: 1763-1777 (1998a); Maloney *et al.*, *Oncology* 12: 63-76 (1998); Leget *et al.*, *Curr. Opin. Oncol.* 10: 548-551 (1998)). In Europe, RITUXAN® has been approved for therapy of relapsed stage III/IV follicular
10 lymphoma (White *et al.*, *Pharm. Sci. Technol. Today* 2: 95-101 (1999)) and is reportedly effective against follicular center cell lymphoma (FCC) (Nguyen *et al.*, *Eur. J. Haematol* 62: 76-82 (1999)). Other disorders treated with RITUXAN® include follicular centre cell lymphoma (FCC), mantle cell lymphoma (MCL), diffuse large cell lymphoma (DLCL), and small
15 lymphocytic lymphoma/chronic lymphocytic leukemia (SLL/CLL) (Nguyen *et al.*, 1999)). Patients with refractory or incurable NHL reportedly have responded to a combination of RITUXAN® and CHOP (*e.g.*, cyclophosphamide, vincristine, prednisone and doxorubicin) therapies (Ohnishi *et al.*, *Gan To Kagaku Ryoho* 25: 2223-8 (1998)). RITUXAN® has exhibited
20 minimal toxicity and significant therapeutic activity in low-grade non-Hodgkin's lymphomas (NHL) in phase I and II clinical studies (Berinstein *et al.*, *Ann. Oncol.* 9: 995-1001 (1998)).

RITUXAN®, which was used alone to treat B-cell NHL at weekly doses of typically 375 mg/M² for four weeks with relapsed or refractory low-grade or
25 follicular NHL, was well tolerated and had significant clinical activity (Piro *et al.*, *Ann. Oncol.* 10: 655-61 (1999); Nguyen *et al.*, (1999); and Coiffier *et al.*, *Blood* 92: 1927-1932 (1998)). However, up to 500 mg/M² of four weekly doses have also been administered during trials using the antibody (Maloney *et al.*, *Blood* 90: 2188-2195 (1997)). RITUXAN® also has been combined with
30 chemotherapeutics, such as CHOP (*e.g.*, cyclophosphamide, doxorubicin, vincristine and prednisone), to treat patients with low-grade or follicular B-cell non-Hodgkin's lymphoma (Czuczman *et al.*, *J. Clin. Oncol.* 17: 268-76 (1999); and McLaughlin *et al.*, (1998a)).

Still further, the use of anti-B7 antibodies for treatment of B cell
35 lymphoma was mentioned in a patent assigned to IDEC Pharmaceuticals Corporation. However, the focus of the patent was the use thereof for treating

5 diseases which immunosuppression is therapeutically beneficial. Examples included allergic, autoimmune and transplant indications. Also mentioned was the use of the discussed anti-B7 antibodies for treatment of B cell lymphoma. (US Patent No. 6,113,198).

B. CD40 and CD40L

10 CD40 is expressed on the cell surface of mature B-cells, as well as on leukemic and lymphocytic B-cells, and on Hodgkin's and Reed-Sternberg (RS) cells of Hodgkin's Disease (HD) (Valle *et al.*, *Eur. J. Immunol.* 19: 1463-1467 (1989); and Gruss *et al.*, *Leuk. Lymphoma* 24: 393-422 (1997)). CD40 is a B-cell receptor leading to activation and survival of normal and malignant B-cells,
15 such as non-Hodgkin's follicular lymphoma (Johnson *et al.*, *Blood* 82: 1848-1857 (1993); and Metkar *et al.*, *Cancer Immunol. Immunother.* 47: 104 (1998)). Signaling through the CD40 receptor protects immature B-cells and B-cell lymphomas from IgM- or Fas-induced apoptosis (Wang *et al.*, *J. Immunology* 155: 3722-3725 (1995)). Similarly, mantle cell lymphoma cells have a high
20 level of CD40, and the addition of exogenous CD40L enhanced their survival and rescued them from fludarabin-induced apoptosis (Clodi *et al.*, *Brit. J. Haematol.* 103: 217-219 (1998)). In contrast, others have reported that CD40 stimulation may inhibit neoplastic B-cell growth both *in vitro* (Funakoshi *et al.*, *Blood* 83: 2787-2794 (1994)) and *in vivo* (Murphy *et al.*, *Blood* 86: 1946-1953
25 (1995)).

Anti-CD40 antibodies (see U.S. Patent Nos. 5,874,082 and 5,667,165) administered to mice increased the survival of mice with human B-cell lymphomas (Funakoshi *et al.*, (1994); and Tutt *et al.*, *J. Immunol.* 161: 3176-3185 (1998)). Methods of treating neoplasms, including B-cell lymphomas and
30 EBV-induced lymphomas using anti-CD40 antibodies mimicking the effect of CD40L and thereby delivering a death signal, are described in U.S. Patent No. 5,674,492 (1997), which is herein incorporated by reference in its entirety. CD40 signaling has also been associated with a synergistic interaction with CD20 (Ledbetter *et al.*, *Circ. Shock* 44: 67-72 (1994)). Additional references
35 describing preparation and use of anti-CD40 antibodies include U.S. Patent Nos. 5,874,085 (1999), 5,874,082 (1999), 5,801,227 (1998), 5,674,492 (1997)

5 and 5,667,165 (1997), which are incorporated herein by reference in their entirety.

A CD40 ligand, gp39 (also called CD40 ligand, CD40L or CD154), is expressed on activated, but not resting, CD4⁺ Th cells (Spriggs *et al.*, *J. Exp. Med.* 176: 1543-1550 (1992); Lane *et al.*, *Eur. J. Immunol.* 22: 2573-2578 (1992); and Roy *et al.*, *J. Immunol.* 151: 1-14 (1993)). Both CD40 and CD40L have been cloned and characterized (Stamenkovi *et al.*, *EMBO J.* 8: 1403-1410 (1989); Armitage *et al.*, *Nature* 357: 80-82 (1992); Lederman *et al.*, *J. Exp. Med.* 175: 1091-1101 (1992); and Hollenbaugh *et al.*, *EMBO J.* 11: 4313-4321 (1992)). Human CD40L is also described in U.S. Patent No. 5,945,513. Cells 15 transfected with the CD40L gene and expressing the CD40L protein on their surface can trigger B-cell proliferation, and together with other stimulatory signals, can induce antibody production (Armitage *et al.*, (1992); and U.S. Patent No. 5,945,513). CD40L may play an important role in the cell contact-dependent interaction of tumor B-cells (CD40⁺) within the neoplastic follicles 20 or Reed-Sternberg cells (CD40⁺) in Hodgkin's Disease areas (Carbone *et al.*, *Am. J. Pathol.* 147: 912-922 (1995)). Anti-CD40L monoclonal antibodies reportedly have been effectively used to inhibit the induction of murine AIDS (MAIDS) in LP-BM5-infected mice (Green *et al.*, *Virology* 241: 260-268 (1998)). However, the mechanism of CD40L-CD40 signaling leading to 25 survival versus cell death responses of malignant B-cells is unclear. For example, in follicular lymphoma cells, down-regulation of a apoptosis inducing TRAIL molecule (APO-2L) (Ribeiro *et al.*, *British J. Haematol.* 103: 684-689 (1998)) and over expression of BCL-2, and in the case of B-CLL, down-regulation of CD95 (Fas/APO-1) (Laytragoon-Lewin *et al.*, *Eur. J. Haematol.* 30 61: 266-271 (1998)) have been proposed as mechanisms of survival. In contrast, evidence exists in follicular lymphoma, that CD40 activation leads to up-regulation of TNF (Worm *et al.*, *International Immunol.* 6: 1883-1890 (1994)) CD95 molecules (Plumas *et al.*, *Blood* 91: 2875-2885 (1998)).

Anti-CD40 antibodies have also been prepared to prevent or treat 35 antibody-mediated diseases, such as allergies and autoimmune disorders as described in U.S. Patent No. 5,874,082 (1999). Anti-CD40 antibodies

5 reportedly have been effectively combined with anti-CD20 antibodies yielding an additive effect in inhibiting growth of non-Hodgkin's B-cell lymphomas in cell culture (Benoit *et al.*, *Immunopharmacology* 35: 129-139 (1996)). *In vivo* studies in mice purportedly demonstrated that anti-CD20 antibodies were more efficacious than anti-CD40 antibodies administered individually in promoting
10 the survival of mice bearing some, but not all, lymphoma lines (Funakoshi *et al.*, *J. Immunother. Emphasis Tumor Immunol.* 19: 93-101 (1996)). Anti-CD19 antibodies are reportedly also effective *in vivo* in the treatment of two syngeneic mouse B-cell lymphomas, BCL1 and A31 (Tutt *et al.* (1998)). Antibodies to CD40L have also been described for use to treat disorders associated with B-
15 cell activation (European Patent No. 555,880 (1993)). Anti-CD40L antibodies include monoclonal antibodies 3E4, 2H5, 2H8, 4D9-8, 4D9-9, 24-31, 24-43, 89-76 and 89-79, as described in U.S. Patent No. 5,747,037 (1998), and anti-CD40L antibodies described in U.S. Patent No. 5,876,718 (1999) used to treat graft-versus-host-disease.

20 C. Anti-CD22 Antibodies

The synthesis of monoclonal antibodies against CD22 and their use in therapeutic regimens has also been reported. CD22 is a B-cell-specific molecule involved in B-cell adhesion that may function in homotypic or heterotypic interactions (Stamenkovic *et al.*, *Nature* 344:74 (1990); Wilson *et al.*,
25 *J. Exp. Med.* 173:137 (1991); Stamenkovic *et al.*, *Cell* 66:1133 (1991)). The CD22 protein is expressed in the cytoplasm of progenitor B and pre-B-cells (Dorken *et al.*, *J. Immunol.* 136:4470 (1986); Dorken *et al.*, "Expression of cytoplasmic CD22 in B-cell ontogeny. In Leukocyte Typing III, White Cell Differentiation Antigens. McMichael *et al.*, eds., Oxford University Press,
30 Oxford, p. 474 (1987); Schwarting *et al.*, *Blood* 65:974 (1985); Mason *et al.*, *Blood* 69:836 (1987)), but is found only on the surface of mature B-cells, being present at the same time as surface IgD (Dorken *et al.*, *J. Immunol.* 136:4470 (1986)). CD22 expression increases following activation and disappears with further differentiation (Wilson *et al.*, *J. Exp. Med.* 173:137 (1991); Dorken *et al.*,
35 *J. Immunol.* 136:4470 (1986)). In lymphoid tissues, CD22 is expressed by follicular mantle and marginal zone B-cells but only weakly by germinal center

5 B-cells (Dorken et al, *J. Immunol.* 136:4470 (1986); Ling et al, "B-cell and
plasma antigens: new and previously defined clusters" In *Leukocyte Typing III.*
White Cell Differentiation Antigens, McMichael et al, eds., Oxford University
Press, Oxford, p. 302 (1987)). However, in situ hybridization reveals the
strongest expression of CD22 mRNA within the germinal center and weaker
10 expression within the mantle zone (Wilson et al, *J. Exp. Med.* 173:137 (1991)).
CD22 is speculated to be involved in the regulation of B-cell activation since
the binding of CD22 mAb to B-cells *in vitro* has been found to augment both
the increase in intracellular free calcium and the proliferation induced after
cross-linking of surface Ig (Pezzutto et al, *J. Immunol.* 138:98 (1987); Pezzutto
15 et al, *J. Immunol.* 140:1791 (1988)). Other studies have determined, however,
that the augmentation of anti-Ig induced proliferation is modest (Dorken et al, *J.*
Immunol. 136:4470 (1986)). CD22 is constitutively phosphorylated, but the
level of phosphorylation is augmented after treatment of cells with PMA (Boue
et al, *J. Immunol.* 140:192 (1988)). Furthermore, a soluble form of CD22
20 inhibits the CD3-mediated activation of human T-cells, suggesting CD22 may
be important in T-cell - B-cell interactions (Stamenkovic et al, *Cell* 66:1133
(1991)).

Ligands that specifically bind the CD22 receptor have been reported to
have potential application in the treatment of various diseases, especially B-cell
25 lymphomas and autoimmune diseases. In particular, the use of labeled and non-
labeled anti-CD22 antibodies for treatment of such diseases has been reported.

For example, Tedder et al, U.S. patent 5,484,892, that purportedly bind
CD22 with high affinity and block the interaction of CD22 with other ligands.
These monoclonal antibodies are disclosed to be useful in treating autoimmune
30 diseases such as glomerulonephritis, Goodpasture's syndrome, necrotizing
vasculitis, lymphadenitis, periarteritis nodosa, systemic lupus erythematosus,
arthritis, thrombocytopenia purpura, agranulocytosis, autoimmune hemolytic
anemias, and for inhibiting immune reactions against foreign antigens such as
fetal antigens during pregnancy, myasthenia gravis, insulin-resistant diabetes,
35 Graves' disease and allergic responses.

5 Also, Leung et al, U. S. Patent 5,789,557, disclose chimeric and humanized anti-CD22 monoclonal antibodies produced by CDR grafting and the use thereof in conjugated and unconjugated form for therapy and diagnosis of B-cell lymphomas and leukemias. The reference discloses especially such antibodies conjugated to cytotoxic agents, such as chemotherapeutic drugs,
10 toxins, heavy metals and radionuclides. (See U.S. Patent 5,789,554, issued August 4, 1998, to Leung et al, and assigned to Immunomedics.)

Further, PCT applications WO 98/42378, WO 00/20864, and WO 98/41641 describe monoclonal antibodies, conjugates and fragments specific to CD22 and therapeutic use thereof, especially for treating B-cell related diseases.

15 Also, the use of anti-CD22 antibodies for treatment of autoimmune diseases and cancer has been suggested. See, e.g., U.S. Patent 5,443,953, issued August 22, 1995 to Hansen et al and assigned to Immunomedics Inc. that purports to describe anti-CD22 immunoconjugates for diagnosis and therapy, especially for treatment of viral and bacterial infectious diseases, cardiovascular
20 disease, autoimmune diseases, and cancer, and U.S. Patent 5,484,892, issued January 16, 1998 to Tedder et al and assigned to Dana-Farber Cancer institute, Inc. that purports to describe various monoclonal antibodies directed against CD22, for treatment of diseases wherein retardation or blocking of CD22 adhesive function is therapeutically beneficial, particularly autoimmune
25 diseases.) These references suggest that an anti-CD22 antibody or fragment may be directly or indirectly conjugated to a desired effector moiety, e.g., a label that may be detected, such as an enzyme, fluorophore, radionuclide, electron transfer agent during an *in vitro* immunoassay or *in vivo* imaging, or a therapeutic effector moiety, e.g., a toxin, drug or radioisotope.

30 Further, an anti-human CD22 monoclonal antibody of the IgG1 isotype is commercially available from Leinco Technologies, and reportedly is useful for treatment of B-cell lymphomas and leukemias, including hairy cell leukemia. (Campana, D. et al, *J. Immunol.* 134:1524 (1985)). Still further, Dorken et al, *J. Immunol.* 150:4719 (1993) and Engel et al, *J. Immunol.*
35 150:4519 (1993) both describe monoclonal antibodies specific to CD22.

5 D. Anti-CD19 Antibodies

Also, the use of anti-CD19 antibodies and fragments thereof for treating lymphoma has been reported in the literature. For example, U. S. Patent 5,686,072, issued November 11, 1997, to Uhr et al, and assigned to the University of Texas, discloses the use of anti-CD19 and anti-CD22 antibodies and immunotoxins for treatment of leukemia lymphomas. This patent is
10 incorporated by reference in its entirety herein.

Further, the use of anti-CD19 antibodies for classifying the status and prognosis of leukemias has been reported.

Thus, based on the foregoing, it is clear that numerous antibodies have
15 been reported to possess therapeutic potential for treatment of B cell malignancies. Notwithstanding this fact, it is an object of the invention to provide novel antibody regimens for treatment of B cell lymphoma.

Brief Description and Objects of the Invention

20 Toward that end, it is an object of the invention to provide a novel improved antibody therapy for treatment of B cell malignancies, including Hodgkin's and non-Hodgkin's lymphoma of any grade.

More specifically, it is an object of the invention to provide a novel antibody regimen for treatment of a B cell malignancy involving the
25 administration of at least one B cell depleting antibody and at least one immunoregulatory or immunomodulatory antibody. -

Even more specifically, it is an object of the invention to provide a novel antibody therapy for treatment of B cell malignancies that involves the administration of at least one B cell depleting antibody preferably selected from
30 an anti-CD20, anti-CD19, anti-CD22 or anti-CD37 antibody and at least one immunomodulatory antibody preferably selected from an anti-B7, anti-CD23, anti-CD40, anti-CD40L or anti-CD4 antibody.

It is another object of the invention to provide a novel therapeutic regimen for treatment of a B cell malignancy such as non-Hodgkin's lymphoma
35 or chronic lymphocyte leukemia (CLL) by the administration of an antibody to CD20 (preferably RITUXAN®) and an antibody to B7 or CD40L (respectively

5 preferably Primatized anti-B7 antibodies reported in US Patent 6,113,198 to Anderson et al, or humanized anti-CD40L antibody reported in US Patent 6,001,358, assigned to IDEC Pharmaceuticals Corporation).

It is another object of the invention to provide novel compositions and kits for treatment of B cell malignancies, in B cell lymphomas and leukemias,
10 that include at least one immunoregulatory or immunomodulatory antibody and at least one B cell depleting antibody. Preferably, the immunoregulatory or immunomodulatory antibody will comprise an anti-CD40, anti-CD40L or anti-B7 antibody and the B cell depleting antibody will be specific to CD20, CD19, CD22 or CD37. Most preferably, the composition will comprise an anti-CD40L
15 or anti-B7 antibody and an anti-CD20 antibody.

Another object of the invention is to provide a combination therapy for the treatment of a B-cell lymphoma or a B-cell leukemia comprising an anti-CD40L antibody or antibody fragment or CD40L antagonist and at least one of the following (a) a chemotherapeutic agent or a combination of
20 chemotherapeutic agents, (b) radiotherapy, (c) an anti-CD20 antibody or fragment thereof, (d) an anti-CD40 antibody or fragment thereof, (e) an anti-CD19 antibody or fragment thereof; (f) an anti-CD22 antibody or fragment thereof, (g) cytokines, and combinations thereof, where antibodies may be conjugated with a toxin or a radiolabel, or may be engineered with human
25 constant regions as to elicit human antibody effector mechanisms, i.e. resulting in apoptosis or death of targeted cells.

Brief Description of the Figures

Fig. 1. Sensitivity of B-lymphoma cells to adriamycin after 4 hour
30 exposure.

Fig. 2. (Panel A) Anti CD40L (IDEC-131) overrides CD40L mediated resistance to killing by ADM of B-lymphoma cells. (Panel B) Effect of RITUXAN® on normal and sCD40L pre-treated DHL-4 cells.

Fig. 3. (Panel A) Blocking of CD40L mediated cell survival of B-CLL
35 by anti-CD40L antibody (IDEC-131). (Panel B) Blocking of CD40L mediated survival of B-CLL by Rituxan®.

- 5 Fig. 4. FACS analysis comprising HLA-DR expression in CD19⁺ CLL cells cultured with sCD40L and not cultured with sCD40L.

Detailed Description of the Invention

- 10 The present invention provides a novel combination antibody regimen that involves the administration of at least one immunoregulatory or immunomodulatory antibody, e.g., an anti-B7 or anti-CD40 or anti-CD40L antibody and at least one B cell depleting antibody, e.g., an anti-CD20, anti-CD19, anti-CD22 or anti-CD37 antibody having substantial B cell depleting activity.
- 15 It is believed that such combination will afford synergistic results based on the different mechanisms by which the antibodies elicit a therapeutic benefit. In particular, it is theorized that the complementary mechanisms of action will yield a more durable and potent clinical response as it is believed that the B cell depleting antibody will deplete activated B cells which may be resistant to the
- 20 action of immunoregulatory or immunomodulatory antibodies such as anti-B7 or anti-CD40L antibodies. Such activated B cells can otherwise serve as effective antigen presenting cells for T cells as well as antibody producing cells. In the context of B cell malignancies, such activated B cells may include malignant cells which unless eradicated by give rise to new cancer cells and
- 25 tumors.

Prior to discussing the invention, the following definitions are provided:

- The term "antibody" as used herein is intended to include immunoglobulins and fragments thereof which are specifically reactive to the designated protein or peptide thereof. An antibody can include human
- 30 antibodies, primatized antibodies, chimeric antibodies, bispecific antibodies, humanized antibodies, antibodies fused to other proteins or radiolabels, and antibody fragments.

- The term "antibody" herein is used in the broadest sense and specifically covers intact monoclonal antibodies, polyclonal antibodies, multispecific
- 35 antibodies (e.g. bispecific antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

5 “Antibody fragments” comprise a portion of an intact antibody,
preferably
comprising the antigen-binding or variable region thereof. Examples of
antibody
fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear
10 antibodies;
single-chain antibody molecules; and multispecific antibodies formed from
antibody
fragments. Antibody fragments may be isolated using conventional techniques.
For example, F(ab')₂ fragments can be generated by treating antibodies with
15 pepsin. The resulting F(ab')₂ fragment can be treated to reduce disulfide
bridges to produce Fab' fragments.

 “Native antibodies” are usually heterotetrameric glycoproteins of about
150,000 daltons, composed of two identical light (L) chains and two identical
heavy (H) chains. Each light chain is linked to a heavy chain by one covalent
20 disulfide bond, while the number of disulfide linkages varies among the heavy
chains of different immunoglobulin isotypes. Each heavy and light chain also
has regularly spaced intrachain disulfide bridges. Each heavy chain has at one
end a variable domain (VH) followed by a number of constant domains. Each
light chain has a variable domain at one end (VL) and a constant domain at its
25 other end; the constant domain of the light chain is aligned with the first
constant domain of the heavy chain, and the light-chain variable domain is
aligned with the variable domain of the heavy chain. Particular amino acid
residues are believed to form an interface between the light chain and heavy
chain variable domains.

30 The term “variable” refers to the fact that certain portions of the variable
domains differ extensively in sequence among antibodies and are used in the
binding and specificity of each particular antibody for its particular antigen.
However, the variability is not evenly distributed throughout the variable
domains of antibodies. It is concentrated in three segments called hypervariable
35 regions both in the light chain and the heavy chain variable domains. The more
highly conserved portions of variable domains are called the framework regions

5 (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a 13-sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the B-sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from 10 the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the 15 antibody in antibody dependent cellular cytotoxicity (ADCC).

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-binding sites 20 and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of 25 each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower 30 affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody 35 hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂Z

5 antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and
10 lambda, based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, antibodies can be assigned to different classes. There are five major classes of
15 intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

The heavy-chain constant domains that correspond to the different classes of
20 antibodies are called alpha, delta, epsilon, gamma and mu, respectively. Preferably, the heavy-chain constant domains will complete the gamma-1, gamma-2, gamma-3 and gamma-4 constant region. Preferably, these constant domains will also comprise modifications to enhance antibody stability such as the P and E modification disclosed in U.S. Patent No. 6,011,138 incorporated
25 by reference in its entirety herein. The subunit structures and three dimensional configurations of different classes of immunoglobulins are well known.

"Single-chain Fv" or "scFv" antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a
30 polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two
35 antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light chain variable domain (VL) in the same polypeptide

5 chain (VH - VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

10 The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal
15 antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal
20 antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the
25 monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*,
30 *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

By "humanized antibody" is meant an antibody derived from a non-human antibody, typically a murine antibody, that retains or substantially retains the antigen-binding properties of the parent antibody, but which is less
35 immunogenic in humans. This may be achieved by various methods, including (a) grafting the entire non-human variable domains onto human constant regions

5 to generate chimeric antibodies; (b) grafting only the non-human complementarity determining regions (CDRs) into human framework and constant regions with or without retention of critical framework residues; and (c) transplanting the entire non-human variable domains, but "cloaking" them with a human-like section by replacement of surface residues. Such methods
10 are disclosed in Morrison *et al.*, *Proc. Natl. Acad. Sci.* 81: 6851-5 (1984); Morrison *et al.*, *Adv. Immunol.* 44: 65-92 (1988); Verhoeyen *et al.*, *Science* 239: 1534-1536 (1988); Padlan, *Molec. Immun.* 28: 489-498 (1991); and Padlan, *Molec. Immun.* 31: 169-217 (1994), all of which are hereby incorporated by reference in their entirety. Humanized anti-CD40L antibodies can be prepared
15 as described in U.S. Patent Application No. 08/554,840 filed November 7, 1995 also incorporated herein by reference in its entirety.

By "human antibody" is meant an antibody containing entirely human light and heavy chain as well as constant regions, produced by any of the known standard methods.

20 By "primatized antibody" is meant a recombinant antibody which has been engineered to contain the variable heavy and light domains of a monkey (or other primate) antibody, in particular, a cynomolgus monkey antibody, and which contains human constant domain sequences, preferably the human immunoglobulin gamma 1 or gamma 4 constant domain (or PE variant). The
25 preparation of such antibodies is described in Newman *et al.*, *Biotechnology*, 10: 1458-1460 (1992); also in commonly assigned 08/379,072, 08/487,550, or 08/746,361, all of which are incorporated by reference in their entirety herein. These antibodies have been reported to exhibit a high degree of homology to human antibodies, *i.e.*, 85-98%, display human effector functions, have reduced
30 immunogenicity, and may exhibit high affinity to human antigens.

By "antibody fragment" is meant an fragment of an antibody such as Fab, F(ab')₂, Fab' and scFv.

By "chimeric antibody" is meant an antibody containing sequences derived from two different antibodies, which typically are of different species.
35 Most typically, chimeric antibodies comprise human and murine antibody fragments, and generally human constant and murine variable regions.

5 "B Cell Depleting Antibody" therein is an antibody or fragment that upon administration, results in demonstrable B cell depletion. Typically, such antibody will bind to a B cell antigen or B cell marker expressed on the surface of a B cell. Preferably, such antibody, after administration, typically within about several days or less, will result in a depletion of B cell number by about 10 50% or more. In a preferred embodiment, the B cell depleting antibody will be RITUXAN® (a chimeric anti-CD20 antibody) or one having substantially the same or at least 20-50% the cell depleting activity of RITUXAN®.

A "B cell surface marker" or "B cell target" or "B cell antigen" herein is an antigen expressed on the surface of a B cell which can be targeted with an 15 antagonist which binds thereto. Exemplary B cell surface markers include the CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD72, CD73, CD74, CDw75, CDw76, CD77, CDw78, CD79a, CD79b, CD80, CD81, CD82, CD83, CDw84, CD85 and CD86 leukocyte surface markers: The B cell surface marker of particular interest is preferentially expressed on B cells compared to 20 other non-B cell tissues of a mammal and may be expressed on both precursor B cells and mature B cells. In one embodiment, the marker is one, like CD20 or CD19, which is found on B cells throughout differentiation of the lineage from the stem cell stage up to a point just prior to terminal differentiation into plasma cells. The preferred B cell surface markers herein are CD19 and CD20.

25 "Immunoregulatory Antibody" refers to an antibody that elicits an effect on the immune system by a mechanism different from B cell depletion, e.g., by CDL and/or ADCC activity. Examples of such include antibodies that inhibit T cell immunity, B cell immunity, e.g. by inducing tolerance (anti-CD40L, anti-CD40) or other immunosuppressant antibodies, e.g., those that inhibit B7 cell 30 signaling (anti-B7.1, anti-B7.2, anti-CD4, anti-CD23, etc.). In some instances, the immunoregulatory antibody may possess the ability to potentiate apoptosis. Also, an antibody that is normally a B cell depleting antibody can be engineered to become immunoregulatory by substantiating human constant regions as to take advantage of different effector mechanisms.

35 A "B cell surface marker" herein is an antigen expressed on the surface of a B cell which can be targeted with an antagonist which binds thereto.

5 Exemplary B cell surface markers include the CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD72, CD73, CD74, CDw75, CDw76, CD77, CDw78, CD79a, CD79b, CD80 (B7.1), CD81, CD82, CD83, CDw84, CD85 and CD86 (B7.2) leukocyte surface markers. The B cell surface marker of particular interest is preferentially expressed on B cells compared to other
10 non-B cell tissues of a mammal and may be expressed on both precursor B cells and mature B cells. In one embodiment, the marker is one, like CD20 or CD 19, which is found on B cells throughout differentiation of the lineage from the stem cell stage up to a point just prior to terminal differentiation into plasma cells. The preferred B cell surface markers herein are CD 19, CD20, CD23, CD80 and CD86.
15

The "CD20" antigen is a -35 kDa, non-glycosylated phosphoprotein found on the surface of greater than 90% of B cells from peripheral blood or lymphoid organs. CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation. CD20 is present on both normal B
20 cells as well as malignant B cells. Other names for CD20 in the literature include "B-lymphocyte-restricted antigen" and "Bp35". The CD20 antigen is described in Clark *et al. PNAS (USA)* 82:1766(1985).

The "CD19" antigen refers to a -90kDa antigen identified, for example, by the

25 HD237-CD19 or B4 antibody (Kiesel *et al. Leukemia Research II*, 12: 1119 (1987)).

Like CD20, CD19 is found on cells throughout differentiation of the lineage from the stem cell stage up to a point just prior to terminal differentiation into plasma cells. Binding of an antagonist to CD 19 may cause internalization of the
30 CD 19 antigen.

The "CD22" antigen refers to an antigen expressed on B cells, also known as "BL-CAM" and "LybB" that is involved in B cell signaling and an adhesion. (See

35 Nitschke *et al., Curr. Biol.* 7:133 (1997); Stamenkovic *et al., Nature* 345:74 (1990)).

- 5 This antigen is a membrane immunoglobulin-associated antigen that is tyrosine phosphorylated when membrane Ig is ligated. (Engel *et al.*, *J. Etyp. Med.* 181(4):1521 1586 (1995)). The gene encoding this antigen has been cloned, and its Ig domains
10 characterized.

B7 antigen includes the B7.1 (CD80), B7.2 (CD81) and B7.3 antigen, which are transmembrane antigens expressed on B cells. Antibodies which specifically bind B7 antigens, including human B7.1 and B7.2 antigens are known in the art. Preferred B7 antibodies comprise the primatized® B7
15 antibodies disclosed by Anderson *et al.* in U.S. Patent No. 6,113,198, assigned to IDEC Pharmaceuticals Corporation, as well as human and humanized B7 antibodies.

CD23 refers to the low affinity receptor for IgE expressed by B and other cells. In the present invention, CD23 will preferably be human CD23
20 antigen. CD23 antibodies are also known in the art. Most preferably, in the present invention, the CD23 antibody will be a human or chimeric anti-human CD23 antibody comprising human IgG1 or IgG3 constant domains.

A B cell "antagonist" is a molecule which, upon binding to a B cell surface marker, destroys or depletes B cells in a mammal and/or interferes with
25 one or more B cell functions, *e.g.* by reducing or preventing a humoral response elicited by the B cell. The antagonist preferably is able to deplete B cells (*i.e.* reduce circulating B cell levels) in a mammal treated therewith. Such depletion may be achieved via various mechanisms such antibody-dependent
30 cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC), inhibition of B cell proliferation and/or induction of B cell death (*e.g.* via apoptosis). Antagonists included within the scope of the present invention include antibodies, synthetic or native sequence peptides and small molecule antagonists which bind to the B cell marker, optionally conjugated with or fused
35 to a cytotoxic agent.

5 A CD40L antagonist is a molecule that specifically binds CD40L and preferably antagonizes the interaction of CD40L and CD40. Examples thereof include antibodies and antibody fragments that specifically bind CD40L, soluble CD40, soluble CD40 fusion proteins, and small molecules that bind CD40L. The preferred antagonist according to the invention comprises an
10 antibody or antibody fragment specific to CD40.

“Antibody-dependent cell-mediated cytotoxicity” and “ADCC” refer to a cell mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (*e.g.* Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the
15 target cell. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US
20 Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be *assessed in vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998).

25 “Human effector cells” are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural
30 killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, *e.g.* from blood or PBMCs as described herein.

The terms “Fc receptor” or “FcR” are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence
35 human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII

5 subclasses, including allelic variants and alternatively spliced forms of these
receptors. FcγRII receptors include FcγRIIA (an “activating receptor”) and
FcγRUB (an “inhibiting receptor”), which have similar amino acid sequences
that differ primarily in the cytoplasmic domains thereof. Activating receptor
FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM)
10 in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an
immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic
domain. (see review M. in Daeon, *Annu. Rev. Immunol.* 15:203-234 (1997)).
FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92
(1991); Capel et al., *Immunomethods* 4:25-34 (1994); and de Haas et al., *J. Lab.*
15 *Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in
the future, are encompassed by the term “FcR” herein. The term also includes
the neonatal receptor, FcRn, which is responsible for the transfer of maternal
IgGs to the fetus (Guyer et al., *J. Immunol.* 117:587 (1976) and Kim et al., *J.*
Immunol. 24:249 (1994)).

20 “Complement dependent cytotoxicity” or “CDC” refers to the ability of
a
molecule to lyse a target in the presence of complement. The complement
activation
pathway is initiated by the binding of the first component of the complement
system (C1q) to a molecule (e.g. an antibody) complexed with a cognate
25 antigen. To assess
complement activation, a CDC assay, e.g. as described in Gazzano-Santoro et
al., *J.*
Immunol. Methods 202:163 (1996), may be performed.

30 “Growth inhibitory” antagonists are those which prevent or reduce
proliferation of a cell expressing an antigen to which the antagonist binds. For
example, the antagonist may prevent or reduce proliferation of B cells in vitro
and/or in vivo.

Antagonists which “induce apoptosis” are those which induce
35 programmed cell death, e.g. of a B cell, as determined by binding of annexin V,

- 5 fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies).

The term "hypervariable region" when used herein refers to the amino acid

residues of an antibody which are responsible for antigen-binding. The

- 10 hypervariable

region comprises amino acid residues from a "complementarity determining region" or "CDR" (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of*

- 15 *Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk. *Mol. Biol.* 196:901-917 (1987)).

- 20 "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

An antagonist "which binds" an antigen of interest, e.g. a B cell surface marker, is one capable of binding that antigen with sufficient affinity such that the antagonist is useful as a therapeutic agent for targeting a cell, i.e. a B cell, expressing the antigen.

- 25

An "anti-CD20 antibody" herein is an antibody that specifically binds CD20 antigen, preferably human CD20, having measurable B cell depleting activity, preferably having at least about 10% the B cell depleting activity of RITUXAN® (see U.S. Patent No. 5,736,137, incorporated by reference herein in its entirety).

- 30

An "anti-CD22 antibody" herein is an antibody that specifically binds CD22 antigen, preferably human CD22, having measurable B cell depleting activity, preferably having at least about 10% the B cell depleting activity of RITUXAN® (see U.S. Patent No. 5,736,137, incorporated by reference herein

- 35 in its entirety).

5 An "anti-CD19 antibody" herein is an antibody that specifically binds CD19 antigen, preferably human CD19, having measurable B cell depleting activity, preferably having at least about 10% the B cell depleting activity of RITUXAN® (see U.S. Patent No. 5,736,137, incorporated by reference herein in its entirety).

10 An "anti-CD37 antibody" herein is an antibody that specifically binds CD37 antigen, preferably human CD37, having measurable B cell depleting activity, preferably having at least about 10% the B cell depleting activity of RITUXAN® (see U.S. Patent No. 5,736,137, incorporated by reference herein in its entirety).

15 An "anti-B7 antibody" herein is an antibody that specifically binds B7.1, B7.2 or B7.3, most preferably human B7.3, that inhibits B7/CD28 interactions and, which more does not substantially inhibit B7/CTLA-4 interactions, and even more preferably, the particular antibodies described in U.S. Patent 6,113,898, incorporated by reference in its entirety herein. It has recently been
20 shown that these antibodies promote apoptosis. Therefore, they are well suited for anti-neoplastic applications.

 An "anti-CD40L antibody" is an antibody that specifically binds CD40L (also known as CD154, gp39, TBAM), preferably one having agonistic activity. A preferred anti-Cd40L antibody is one having the specificity of a humanized
25 antibody disclosed in U.S. Patent No. 6,011,358 (assigned to IDEC Pharmaceuticals Corporation), incorporated by reference in its entirety herein.

 An "anti-CD4 antibody" is one that specifically binds CD4, preferably human CD4, more preferably a primatized or humanized anti-CD4 antibody.

 An "anti-CD40 antibody" is an antibody that specifically binds CD40,
30 preferably human CD40, such as those disclosed in U.S. Patent 5,874,085, 5,874,082, 5,801,227, 5,674,442, snf 5,667,165, all of which are incorporated by reference herein.

 Preferably, both the B cell depleting antibody and the immunoregulatory antibody will contain human constant domains. Suitable antibodies may
35 include IgG1, IgG2, IgG3 and IgG4 isotypes.

5 Specific examples of antibodies which bind the CD20 antigen include:

“Rituximab” (“RITUXAN®”) (US Patent No. 5,736,137, expressly
incorporated herein by reference); yttrium-[90]-labeled 2B8 murine antibody
“Y2B8” (US Patent No. 5,736,B7, expressly incorporated herein by reference);
murine IgG2a “B1” optionally labeled with 1311, «1311 B1” antibody
10 (BEXXARTM) (US Patent No. 5,595,721, expressly incorporated herein by
reference); murine monoclonal antibody “1F5” (Press *et al. Blood*
69(2):584-591 (1987); and “chimeric 2H7” antibody (US Patent No. 5,677,180,
expressly incorporated herein by reference).

Specific examples of antibodies which bind CD22 include

15 Lymphocide™ reported by Immunomedics, now in clinical trials for
non-Hodgkin’s lymphoma. Examples of antibodies that bind B7 antigen include
the B7 antibody reported U.S. Patent 5,885,577, issued to Linsley et al, the
anti-B7 antibody reported in U.S. Patent 5,869,050, issued in DeBoer et al,
assigned to Chiron Corporation, and the primatized® anti-B7 antibodies
20 disclosed in U.S. Patent 6,113,198 to Anderson et al., all of which are
incorporated by reference in their entirety.

Specific examples of antibodies that bind CD23 are well known and
preferably include the primatized® antibodies specific to human CD23 reported
by Reff *et al.*, in U.S. Patent 6,011,138, issued on July 4, 1999, co-assigned to
25 IDEC Pharmaceuticals Corp. and Seikakagu Corporation of Japan; those
reported by Bonnefoy *et al.*, No. 96 12741; Rector *et al. J. Immunol.*
55:481-488 (1985); Flores-Rumeo *et al. Science* 241:1038-1046 (1993); Sherr
et al. J. Immunol., 142:481-489 (1989); and Pene *et al., PNAS, USA*
85:6820-6824 (1988). Such antibodies are reportedly useful for treatment of
30 allergy, autoimmune diseases, and inflammatory diseases.

The terms “rituximab” or “RITUXAN®” herein refer to the genetically
engineered chimeric murine/human monoclonal antibody directed against the
CD20
antigen and designated “C2B8” in US Patent No. 5,736,B7, expressly
35 incorporated

5 herein by reference. The antibody is an IgG1 kappa immunoglobulin containing murine light and heavy chain variable region sequences and human constant region sequences. Rituximab has a binding affinity for the CD20 antigen of approximately 8.0 nM.

10 An "isolated" antagonist is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antagonist, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred
15 embodiments, the antagonist will be purified (1) to greater than 95% by weight of antagonist as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions
20 using Coomassie blue or, preferably, silver stain. Isolated antagonist includes the antagonist *in situ* within recombinant cells since at least one component of the antagonist's natural environment will not be present. Ordinarily, however, isolated antagonist will be prepared by at least one purification step.

25 "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, *etc.* Preferably, the mammal is human.

30 "Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disease or disorder as well as those in which the disease or disorder is to be prevented. Hence, the mammal may have been diagnosed as having the disease or disorder or may be predisposed or susceptible to the disease.

B Cell Malignancy

35 According to the present invention this includes any B cell malignancy, e.g., B cell lymphomas and leukemias. Preferred examples include Hodgkin's disease (all forms, e.g., relapsed Hodgkin's disease, resistant Hodgkin's disease)

5 non-Hodgkin's lymphomas (low grade, intermediate grade, high grade, and other types). Examples include small lymphocytic/B cell chronic lymphocytic leukemia (SLL/B-CLL), lymphoplasmacytoid lymphoma (LPL), mantle cell lymphoma (MCL), follicular lymphoma (FL), diffuse large cell lymphoma (DLCL), Burkitt's lymphoma (BL), AIDS-related lymphomas, monocytic B
10 cell lymphoma, angioimmunoblastic lymphadenopathy, small lymphocytic, follicular, diffuse large cell, diffuse small cleaved cell, large cell immunoblastic lymphoblastoma, small, non-cleaved, Burkitt's and non-Burkitt's, follicular, predominantly large cell; follicular, predominantly small cleaved cell; and follicular, mixed small cleaved and large cell lymphomas. See, Gaidono et al.,
15 "Lymphomas", IN CANCER: PRINCIPLES & PRACTICE OF ONCOLOGY, Vol. 2: 2131-2145 (DeVita et al., eds., 5th ed. 1997).

Other types of lymphoma classifications include immunocytomal Waldenstrom's MALT-type/monocytoid B cell, mantle cell lymphoma B-
CLL/SLL, diffuse large B-cell lymphoma, follicular lymphoma, and precursor
20 B-LBL.

As noted, B cell malignancies further include especially leukemias such as ALL-L3 (Burkitt's type leukemia), chronic lymphocytic leukemia (CLL), chronic leukocytic leukemia, acute myelogenous leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia,
25 lymphoblastic leukemia, lymphocytic leukemia, monocytic leukemia, myelogenous leukemia, and promyelocytic leukemia and monocytic cell leukemias.

The expression "therapeutically effective amount" refers to an amount of the
30 antagonist which is effective for preventing, ameliorating or treating the B cell malignancy disease in question.

The term "immunosuppressive agent" as used herein for adjunct therapy refers to substances that act to suppress or mask the immune system of the mammal being treated herein. This would include substances that suppress
35 cytokine production, downregulate or suppress self-antigen expression, or mask the MHC antigens. Examples of such agents include

- 5 2-amino-6-aryl-5-substituted pyrimidines (see U.S. Pat. No. 4,665,077, the disclosure of which is incorporated herein by reference), azathioprine; cyclophosphamide; bromocryptine; danazol; dapsone; glutaraldehyde (which masks the MHC antigens, as described in U.S. Pat. No. 4,120,649); anti-idiotypic antibodies for MHC antigens and MHC fragments; cyclosporin A;
- 10 steroids such as glucocorticosteroids, *e.g.*, prednisone, methylprednisolone, and dexamethasone; cytokine or cytokine receptor antagonists including anti-interferon- α , β - or δ -antibodies, anti-tumor necrosis factor- α antibodies, anti-tumor necrosis factor- β antibodies, anti-interleukin-2 antibodies and anti-IL-2 receptor antibodies; anti-LFA-1 antibodies, including anti-CD11a and
- 15 anti-CD18 antibodies; anti-L3T4 antibodies; heterologous anti-lymphocyte globulin; pan-T antibodies, preferably anti-CD3 or anti-CD4/CD4a antibodies; soluble peptide containing a LFA-3 binding domain (WO 90/08187 published 7/26/90), streptolysin; TGF- β ; streptodornase; RNA or DNA from the host; FK506; RS-61443; deoxyspergualin; rapamycin; T-cell receptor (Cohen *et al.*,
- 20 U.S. Pat. No. 5,114,721); T-cell receptor fragments (Offner *et al.*, *Science*, 251: 430-432 (1991); WO 90/11294; laneway, *Nature*, 341: 482 (1989); and WO 91/01133); and T cell receptor antibodies (EP 340,109) such as T10B9.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or

- 25 prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e. g.* At211-1131 1125 Y90 Re 186 Re 188 Sm153 Bi212 p32 and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

- 30 A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and
- 35 methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and

- 5 trimethylolomelamine nitrogen mustards such as chiorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such
- 10 as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins,
- 15 peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine;
- 20 pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofer, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanol, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolic acid; aceglatone;
- 25 aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diazi quone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine;
- 30 PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triazi quone; 2, 2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, NJ) and doxetaxel (Taxotere, Rhone-Poulenc
- 35 Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and

5 carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any
10 of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and antiandrogens such as flutamide, nilutamide,
15 bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide
20 hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic
25 growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF-13; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and
30 TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocytemacrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1a, IL-2, IL-g, IL-4, IL-5, IL-6, IL-7, IL-8,
35 IL-9, IL-11, IL-12, IL-15; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the

5 term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to tumor cells compared to the parent drug and is capable of being enzymatically
10 activated or converted into the more active parent form. See, *e.g.*, Wilman, "Prodrugs in Cancer Chemotherapy" *Biochemical Society Transactions*, 14, pp. 375-382, 615th Meeting Belfast (1986) and Stella *et al.*, "Prodrugs: A Chemical Approach to Targeted Drug Delivery," *Directed Drug Delivery*, Borchardt *et al.*, (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention
15 include, but are not limited to, phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, 13-lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted
20 phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described above.

25 A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the antagonists disclosed herein and, optionally, a chemotherapeutic agent) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

30 The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

35

5 II. Production of Antibodies

The methods and articles of manufacture of the present invention use, or incorporate, an antibody that has immunoregulatory activity, e.g. anti-B7, anti-CD23, anti-CD40L, anti-CD4 or anti-CD40 antibody, and an antibody that binds to a B cell surface marker having B depleting activity, e.g., anti-CD20,
10 anti-CD22, anti-CD19, or anti-CD37 antibody. Accordingly, methods for generating such antibodies will be described herein.

The molecule to be used for production of, or screening for, antigen(s) may be, e.g., a soluble form of the antigen or a portion thereof, containing the desired epitope. Alternatively, or additionally, cells expressing the antigen at
15 their cell surface can be used to generate, or screen for, antagonist(s). Other forms of the B cell surface marker useful for generating antagonists will be apparent to those skilled in the art. Suitable antigen sources for CD40L, CD40, CD19, CD20, CD22, CD23, CD37, CD4 and B7 antigen (e.g., B7.1, B7.2) antigen for producing antibodies according to the invention are well known.
20 Alternatively, peptides can be synthetically prepared based upon the amino acid sequence. For example, with respect to CD40L, this is disclosed in Armitage et al. (1992).

Preferably, the CD40L antibody or anti-CD40L antibody will be the humanized anti-CD40L antibody disclosed in U.S. Patent 6,001,358, issued on
25 June 14, 1999, and assigned to IDEC Pharmaceuticals Corporation.

While a preferred CD40L antagonist is an antibody, antagonists other than antibodies may also be administered. For example, the antagonist may comprise soluble CD40, a CD40 fusion protein or a small molecule antagonist optionally fused to, or conjugated with, a cytotoxic agent (such as those
30 described herein). Libraries of small molecules may be screened against the B cell surface marker of interest herein in order to identify a small molecule which binds to that antigen. The small molecule may further be screened for its antagonistic properties and/or conjugated with a cytotoxic agent.

The antagonist may also be a peptide generated by rational design or by
35 phage

- 5 display (W098/35036 published 13 August 1998), for example. In one embodiment, the molecule of choice may be a "CDR mimic" or antibody analogue designed based on the CDRs of an antibody, for example. While the peptide may be antagonistic by itself, the peptide may optionally be fused to a cytotoxic agent or to an immunoglobulin Fc region (*e.g.*, so as to confer ADCC
10 and/or CDC activity on the peptide).

Exemplary techniques for the production of the antibody antagonists used in accordance with the present invention are described.

(i) *Polyclonal antibodies*

- 15 Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor
20 using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOC I₂, or R¹N=C=NR, where R and R¹ are different alkyl groups.

- Animals are immunized against the antigen, immunogenic conjugates, or
25 derivatives by combining, *e.g.* 100 µg or 5 µg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites.
30 Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also,
35 aggregating agents such as alum are suitably used to enhance the immune response.

5 (ii) *Monoclonal antibodies*

Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are

10 identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the monoclonal antibodies may be made using the hybridoma

15 method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that
20 produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as
25 polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable
30 culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma

5 cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase
(HGPRT or
HPRT), the culture medium for the hybridomas typically will include
hypoxanthine,
aminopterin, and thymidine (HAT medium), which substances prevent the
10 growth of
HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable
high-level production of antibody by the selected antibody-producing cells, and
are sensitive to a medium such as HAT medium. Among these, preferred
15 myeloma cell lines are murine myeloma lines, such as those derived from
MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell
Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653
cells available from the American Type Culture Collection, Manassas, Virginia,
USA. Human myeloma and mouse-human heteromyeloma cell lines also have
20 been described for the production of human monoclonal antibodies (Kozbor, J.
Immunol., 133:300 1 (1984); Brodeur *et al.*, *Monoclonal Antibody Production
Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York,
1987)).

Culture medium in which hybridoma cells are growing is assayed for
25 production of monoclonal antibodies directed against the antigen. Preferably,
the binding specificity of monoclonal antibodies produced by hybridoma cells is
determined by immunoprecipitation or by an *in vitro* binding assay, such as
radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be
30 determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*,
107:220 (1980).

After hybridoma cells are identified that produce antibodies of the
desired specificity, affinity, and/or activity, the clones may be subcloned by
limiting dilution procedures and grown by standard methods (Goding,
35 *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press,
1986)). Suitable culture media for this purpose include, for example, D-MEM or

- 5 RPML-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein
10 A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light
15 chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal
20 antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Pluckthun, *Immunol. Revs.*, 130:151-188 (1992).

Another method of generating specific antibodies, or antibody
25 fragments, reactive against a CD40L, CD19, CD22, CD20, or CD40 protein or peptide (*e.g.*, such as the gp39 fusion protein described in U.S. Patent No. 5,945,513) is to screen expression libraries encoding immunoglobulin genes, or portions thereof, expressed in bacteria with a CD40L, CD19, CD20, or CD22 protein or peptide. For example, complete Fab fragments, V_H regions and Fv
30 regions can be expressed in bacteria using phage expression libraries. See for example, Ward *et al.*, *Nature* 341: 544-546 (1989); Huse *et al.*, *Science* 246: 1275-1281 (1989); and McCafferty *et al.*, *Nature* 348: 552-554 (1990). Screening such libraries with, for example, a CD40L, CD22, CD19, or CD20 peptide, can identify immunoglobulin fragments reactive with CD40L, CD22,
35 CD19, or CD20. Alternatively, the SCID-hu mouse (available from Genpharm) can be used to produce antibodies or fragments thereof.

5 In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554(1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using
10 phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these
15 techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

Methodologies for producing monoclonal antibodies (MAb) directed against CD40L, including human CD40L and mouse CD40L, and suitable monoclonal antibodies for use in the methods of the invention, are described in
20 PCT Patent Application No. WO 95/06666 entitled "Anti-gp39 Antibodies and Uses Therefor," the teachings of which are incorporated herein by reference in their entirety. Particularly preferred anti-human CD40L antibodies of the invention are MAbs 24-31 and 89-76, produced respectively by hybridomas 24-31 and 89-76. The 89-76 and 24-31 hybridomas, producing the 89-76 and 24-
25 31 antibodies, respectively, were deposited under the provisions of the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, VA 20110-2209, on Sept. 2, 1994. The 89-76 hybridoma was assigned ATCC Accession Number HB11713 and the 24-31 hybridoma was assigned ATCC Accession Number HB11712.

30 The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a
35 non-immunoglobulin polypeptide.

5 Typically, such non-immunoglobulin polypeptides are substituted for the
constant domains of an antibody, or they are substituted for the variable
domains of one antigencombining site of an antibody to create a chimeric
bivalent antibody comprising one antigen-combining site having specificity for
an antigen and another antigen-combining site having specificity for a different
10 antigen.

(iii) *Humanized antibodies*

Methods for humanizing non-human antibodies have been described in
the art.

15 Preferably, a humanized antibody has one or more amino acid residues
introduced into it from a source which is non-human. These non-human amino
acid residues are often referred to as "import" residues, which are typically
taken from an "import" variable domain. Humanization can be essentially
performed following the method of Winter and co-workers (Jones *et al.*, *Nature*,
20 321:522-525 (1986); Reichmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeven
et al., *Science*, 239:1534-1536 (1988)), by substituting hypervariable region
sequences for the corresponding sequences of a human antibody. Accordingly,
such "humanized" antibodies are chimeric antibodies (U.S. Patent No.
4,816,567) wherein substantially less than an intact human variable domain has
25 been substituted by the corresponding sequence from a non-human species. In
practice, humanized antibodies are typically human antibodies in which some
hypervariable region residues and possibly some FR residues are substituted by
residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used
30 in making the humanized antibodies is very important to reduce antigenicity.
According to the so called "best-fit" method, the sequence of the variable
domain of a rodent antibody is screened against the entire library of known
human variable-domain sequences. The human sequence which is closest to that
of the rodent is then accepted as the human framework region (FR) for the
35 humanized antibody (Suns *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*,
J. Mol. Biol., 196:901 (1987)). Another method uses a particular framework

5 region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of
10 high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are
15 commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i. e., the analysis of
20 residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in
25 influencing antigen binding.

(iv) *Primatized Antibodies*

Another highly efficient means for generating recombinant antibodies is disclosed by Newman, *Biotechnology*, 10: 1455-1460 (1992). More
30 particularly, this technique results in the generation of primatized antibodies which contain monkey variable domains and human constant sequences. This reference is incorporated by reference in its entirety herein. Moreover, this technique is also described in commonly assigned U.S. Application No. 08/379,072, filed on January 25, 1995, which is a continuation of U.S. Serial
35 No. 07/912,292, filed July 10, 1992, which is a continuation-in-part of U.S. Serial No. 07/856,281, filed March 23, 1992, which is finally a continuation-in-

- 5 part of U.S. Serial No. 07/735,064, filed July 25, 1991. 08/379,072 and the parent application thereof all of which are incorporated by reference in their entirety herein.

10 This technique modifies antibodies such that they are not antigenically rejected upon administration in humans. This technique relies on immunization of cynomolgus monkeys with human antigens or receptors. This technique was developed to create high affinity monoclonal antibodies directed to human cell surface antigens.

15 Identification of macaque antibodies to human CD40L, CD20, CD22, CD40 or CD19 by screening of phage display libraries or monkey heterohybridomas obtained using B lymphocytes from CD40L, CD20, CD22, CD40, or CD19 immunized monkeys can be performed using the methods described in commonly assigned U.S. Application No. 08/487,550, filed June 7, 1995, incorporated by reference in its entirety herein.

20 Antibodies generated using the methods described in these applications have previously been reported to display human effector function, have reduced immunogenicity, and long serum half-life. The technology relies on the fact that despite the fact that cynomolgus monkeys are phylogenetically similar to humans, they still recognize many human proteins as foreign and therefore mount an immune response. Moreover, because the cynomolgus monkeys are phylogenetically close to humans, the antibodies generated in these monkeys have been discovered to have a high degree of amino acid homology to those produced in humans. Indeed, after sequencing macaque immunoglobulin light and heavy chain variable region genes, it was found that the sequence of each gene family was 85-98% homologous to its human counterpart (Newman *et al.*, 25 1992). The first antibody generated in this way, an anti-CD4 antibody, was 91-92% homologous to the consensus sequence of human immunoglobulin framework regions (Newman *et al.*, 1992).

35 As described above, the present invention relates, in part, to the use of monoclonal antibodies or primatized forms thereof which are specific to human CD40L antigen and which are capable of inhibiting CD40 signaling or inhibiting CD40/CD40L interaction. Blocking of the primary activation site

5 between CD40 and CD40L with the identified antibodies (or therapeutically effective fragments thereof), while allowing the combined antagonistic effect on positive co-stimulation with an agnostic effect on negative signaling will be a useful therapeutic approach for intervening in relapsed forms of malignancy, especially B-cell lymphomas and leukemias. The functional activity of the
10 identified antibodies is defined by blocking the signals of CD40 permitting it to survive and avoid IgM- or Fas-induced apoptosis.

Manufacture of monoclonal antibodies which specifically bind human
CD40L, as well as primatized antibodies derived therefrom can be performed using the methods described in U.S. Patent Nos. 6,001,358 or 5,750,105, both
15 assigned to IDEC Pharmaceuticals Corporation, or other known methods. Preferably, such antibodies will possess high affinity to CD40L and therefore may be used as immunosuppressants which inhibit the CD40L/CD40 pathway. Similar techniques will yield monkey antibodies specific to CD20, CD19, CD22 or CD40.

20 Preparation of monkey monoclonal antibodies will preferably be effected by screening of phage display libraries or by preparation of monkey heterohybridomas using B lymphocytes obtained from CD40L (e.g., human CD40L) immunized monkeys. The human CD40 can also be from the fusion protein described in U.S. Patent No. 5,945,513.

25 As noted, the first method for generating anti-CD40L, CD19, CD20, CD22 or CD40 antibodies involves recombinant phage display technology. This technique is generally described *supra*.

Essentially, this will comprise synthesis of recombinant immunoglobulin libraries against the target, i.e., CD19, CD22, CD20, CD40, or CD40L antigen
30 displayed on the surface of filamentous phage and selection of phage which secrete antibodies having high affinity to CD40L antigen. As noted *supra*, preferably antibodies will be selected which bind to both human CD40L and CD40. To effect such methodology, the present inventors have created a unique library for monkey libraries which reduces the possibility of recombination and
35 improves stability.

5 Essentially, to adopt phage display for use with macaque libraries, this vector contains specific primers for PCR amplifying monkey immunoglobulin genes. These primers are based on macaque sequences obtained while developing the primatized technology and databases containing human sequences.

10 Suitable primers are disclosed in commonly assigned 08/379,072, incorporated by reference herein.

 The second method involves the immunization of monkeys, *i.e.*, macaques, against the desired antigen target, *i.e.*, human CD19, CD20, CD22, CD40 or CD40L. The inherent advantage of macaques for generation of
15 monoclonal antibodies is discussed *supra*. In particular, such monkeys, *i.e.*, cynomolgus monkeys, may be immunized against human antigens or receptors. Moreover, the resultant antibodies may be used to make primatized antibodies according to the methodology of Newman *et al.*, (1992), and Newman *et al.*, commonly assigned U.S. Serial No. 08/379,072, filed January 25, 1995, which
20 are incorporated by reference in their entirety.

 The significant advantage of antibodies obtained from cynomolgus monkeys is that these monkeys recognize many human proteins as foreign and thereby provide for the formation of antibodies, some with high affinity to desired human antigens, *e.g.*, human surface proteins and cell receptors.
25 Moreover, because they are phylogenetically close to humans, the resultant antibodies exhibit a high degree of amino acid homology to those produced in humans. As noted above, after sequencing macaque immunoglobulin light and heavy variable region genes, it was found that the sequence of each gene family was 85-88% homologous to its human counterpart (Newman *et al.*, 1992).

30 Essentially, cynomolgus macaque monkeys are administered human, CD19, CD20, CD22, CD40, or CD40L antigen, B cells are isolated therefrom, *e.g.*, lymph node biopsies are taken from the animals, and B lymphocytes are then fused with KH6/B5 (mouse x human) heteromyeloma cells using polyethylene glycol (PEG). Heterohybridomas secreting antibodies which bind
35 human CD40L antigen are then identified.

5 In the case of antibodies which bind to CD40L or CD40, it is desirable that they do so in a manner which interrupts or regulates CD40 signaling because such antibodies potentially may be used to inhibit the interaction of CD40L with CD40, with their counter-receptors. If antibodies can be developed against more than one epitope on CD40L or CD40, and the
10 antibodies are utilized together, their combined activity may potentially provide synergistic effects.

The disclosed invention involves the use of an animal which is primed to produce a particular antibody (*e.g.*, primates, such as orangutan, baboons, macaque, and cynomolgus monkeys). Other animals which may be used to
15 raise antibodies to human CD40L include, but are not limited to, the following: mice, rats, guinea pigs, hamsters, monkeys, pigs, goats and rabbits.

Cell lines which express antibodies which specifically bind to human CD40L antigen are then used to clone variable domain sequences for the manufacture of primatized antibodies essentially as described in Newman *et al.*,
20 (1992) and Newman *et al.*, U.S. Serial No. 379,072, filed January 25, 1995, both of which are incorporated by reference herein. Essentially, this entails extraction of RNA therefrom, conversion to cDNA, and amplification thereof by PCR using Ig specific primers. Suitable primers are described in Newman *et al.*, 1992, and in U.S. Serial No. 379,072. Similar techniques will yield cell lines
25 that express antibodies specific to CD40, CD19, CD20, or CD22.

Cell lines which express antibodies which specifically bind to human CD40L antigen are then used to clone variable domain sequences for the manufacture of primatized antibodies essentially as described in Newman *et al.*, (1992) and Newman *et al.*, U.S. Serial No. 379,072, filed January 25, 1995,
30 both of which are incorporated by reference herein. Essentially, this entails extraction of RNA therefrom, conversion to cDNA, and amplification thereof by PCR using Ig specific primers. Suitable primers are described in Newman *et al.*, 1992, and in U.S. Serial No. 379,072. Similar techniques will yield cell lines that express antibodies specific to CD40, CD19, CD20, or CD22.

35 Cell lines which express antibodies which specifically bind to human CD40L antigen are then used to clone variable domain sequences for the

5 manufacture of primatized antibodies essentially as described in Newman *et al.*,
(1992) and Newman *et al.*, U.S. Serial No. 379,072, filed January 25, 1995,
both of which are incorporated by reference herein. Essentially, this entails
extraction of RNA therefrom, conversion to cDNA, and amplification thereof
by PCR using Ig specific primers. Suitable primers are described in Newman *et al.*
10 *et al.*, 1992, and in U.S. Serial No. 379,072. Similar techniques will yield cell lines
that express antibodies specific to CD40, CD19, CD20, or CD22.

The cloned monkey variable genes are then inserted into an expression
vector which contains human heavy and light chain constant region genes.
Preferably, this is effected using a proprietary expression vector of IDEC, Inc.,
15 referred to as NEOSPLA. This vector contains the cytomegalovirus
promoter/enhancer, the mouse beta globin major promoter, the SV40 origin of
replication, the bovine growth hormone polyadenylation sequence, neomycin
phosphotransferase exon 1 and exon 2, human immunoglobulin kappa or
lambda constant region, the dihydrofolate reductase gene, the human
20 immunoglobulin gamma 1 or gamma 4 PE constant region and leader sequence.
This vector has been found to result in very high level expression of primatized
antibodies upon incorporation of monkey variable region genes, transfection in
CHO cells, followed by selection in G418 containing medium and methotrexate
amplification.

25 For example, this expression system has been previously disclosed to
result in primatized antibodies having high avidity ($K_d \leq 10^{-10}$ M) against CD4
and other human cell surface receptors. Moreover, the antibodies have been
found to exhibit the same affinity, specificity and functional activity as the
original monkey antibody. This vector system is substantially disclosed in
30 commonly assigned U.S. Serial No. 379,072, incorporated by reference herein
as well as U.S. Serial No. 08/149,099, filed on November 3, 1993, also
incorporated by reference in its entirety herein. This system provides for high
expression levels, *i.e.*, > 30 pg/cell/day. Of course, the same methods can be
used to produce cell lines that produce antibodies specific to CD19, CD20,
35 CD22, or CD40.

5 (v) *Human antibodies*

As an alternative to humanization, human antibodies can be generated.

For

example, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies

10 in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region PH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ line mutant mice will result
15 in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits *et al.*, *Proc. Mad. Acad. Sci. USA*, 90:255-1 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in immuno.*, 7:33 (1993); and US Patent Nos. 5,591,669, 5,589,369 and 5,545,807.

Alternatively, phage display technology (McCafferty *et al.*, *Nature*
20 348:552-553 (1990)) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional
25 antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B cell. Phage display can be
30 performed in a variety of formats; for their review see, e.g., Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology* 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson *et al.*, *Nature*, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes
35 derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse

5 array of antigens (including self antigens) can be isolated essentially following the techniques described by Marks *et al.*, *J.Mol. Biol.* 222:581-597 (1991), or Griffith *et al.*, *EMBO J.* 12:725-734 (1993). See, also, US Patent Nos. 5,565,332 and 5,573,905.

10 Human antibodies may also be generated by in *vitro* activated B cells (see US Patents 20 5,567,610 and 5,229,275). A preferred means of generating human antibodies using SCID mice is disclosed in commonly-owned, co-pending applications.

15 (vi) *Antibody fragments*

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81
20 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10: 163-167 (1992)).
25 According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; US Patent No. 5,571,894; and US Patent No. 5,587,458. The antibody fragment
30 may also be a "linear antibody", *e.g.*, as described in US Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

(vii) *Bispecific antibodies*

Bispecific antibodies are antibodies that have binding specificities for at
35 least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the B cell surface marker. Other such antibodies may bind

5 a first B cell marker and further bind a second B cell surface marker.

Alternatively, an anti-B cell marker binding arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD 16) so as to focus cellular defense mechanisms to the B cell. Bispecific antibodies may also be used to localize cytotoxic agents to the B cell. These antibodies possess a B cell marker-binding arm and an arm which binds the cytotoxic agent (*e.g.* saporin, anti-interferon-α, vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten).

10 Bispecific antibodies can be prepared as full length antibodies or antibody fragments (*e.g.* F(ab)₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art.

Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low.

20 Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin

30 constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for

35

5 great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in
10 equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain
15 pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO
20 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach described in US Patent No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize
25 the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from
30 the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chains) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the
35 heterodimer over other unwanted end-products such as homodimers.

5 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for
10 treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

15 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229:81(1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate
20 F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with
25 mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

 Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies.
30 Shalaby *et al.*, *J.Exp. Med.*, 175:2 17-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T
35 cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

5 Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two
10 different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative
15 mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment,
20 thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147:
25 60(1991).

III. Conjugates and Other Modifications of the Antagonist

The antibodies used in the methods or included in the articles of manufacture herein are optionally conjugated to a cytotoxic agent.

30 Chemotherapeutic agents useful in the generation of such antibody-cytotoxic agent conjugates have been described above.

Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, a maytansine (US Patent No. 5,208,020), a trichothene, and
35 CC 1065 are also contemplated herein. In one preferred embodiment of the invention, the antagonist is conjugated to one or more maytansine molecules

5 (e.g. about 1 to about 10 maytansine molecules per antagonist molecule).

Maytansine may, for example, be converted to May SS-Me which may be reduced to May-SH3 and reacted with modified antagonist (Charm *et al. Cancer Research* 52:127-131(1992)) to generate a maytansinoid-antagonist conjugate.

Alternatively, the antibody may be conjugated to one or more

10 calicheamicin

molecules. The calicheamicin family of antibiotics are capable of producing double

stranded DNA breaks at sub-picomolar concentrations. Structural analogues of calicheamicin which may be used include, but are not limited to, γ_1^I , α_2^I , α_3^I ,

15 N-acetyl- γ_1^I , PSAG and O_1^I (Hinman *et al. Cancer Research* 53:3336-3342 (1993) and Lode *et al. Cancer Research* 58: 2925-2928 (1998)).

Enzymatically active toxins and fragments thereof which can be used include

diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A

20 chain

(from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates antibody conjugated with a compound with nucleolytic activity (e.g. a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

30 A variety of radioactive isotopes are available for the production of radioconjugated antagonists. Examples include At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} and radioactive isotopes of Lu.

Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as

35 N-succinimidyl-3-(2-pyridyldithiol)

- 5 propionate (SPDP), succinimidyl-4-(N-maleimidomethyl)
cyclohexane-1-carboxylate,
iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl
adipimide HCL), active esters (such as disuccinimidyl suberate), aldehydes
(such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl)
10 hexanediamine), bis-diazonium derivatives (such as
bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene
2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,
4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as
described in Vitetta *et al. Science* 238: 1098 (1987). Carbon-14-labeled 1
15 isothiocyanatobenzyl-3- methyl-diethylene triamine-pentaacetic acid
(MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide
to the antagonist. See W094/11026. The linker may be a "cleavable linker"
facilitating release of the cytotoxic drug in the cell. For example, an acid-labile
linker, peptidase-sensitive linker, dimethyl linker or disulfide-containing linker
20 (Charm *et al. Cancer Research* 52:127-131 (1992)) may be used.

Alternatively, a fusion protein comprising the antibody and cytotoxic
agent may be made, *e.g.* by recombinant techniques or peptide synthesis.

- In yet another embodiment, the antibody may be conjugated to a
"receptor" (such streptavidin) for utilization in tumor pretargeting wherein the
25 antagonist-receptor conjugate is administered to the patient, followed by
removal of unbound conjugate from the circulation using a clearing agent and
then administration of a "ligand" (*e.g.* avidin) which is conjugated to a cytotoxic
agent (*e.g.* a radionucleotide).

- The antibodies of the present invention may also be conjugated with a
30 prodrug activating enzyme which converts a prodrug (*e.g.* a peptidyl
chemotherapeutic agent, see W081/01145) to an active anti-cancer drug. See,
for example, WO 88/07378 and U.S. Patent No. 4,975,278.

- The enzyme component of such conjugates includes any enzyme
capable of acting on a prodrug in such a way so as to convert it into its more
35 active, cytotoxic form.

5 Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, fluorouracil; proteases, 10 such as serratiopeptidase, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase 15 useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with 20 enzymatic activity, also known in the art as "abzymes", can be used to convert the prodrugs of the invention into free active drugs (see, *e.g.*, Massey, *Nature* 328:457-458 (1987)). Antagonist-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes of this invention can be covalently bound to the antagonist 25 by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antagonist of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed 30 using recombinant DNA techniques well known in the art (see, *e.g.*, Neuberger *et al.*, *Nature*, 312:604-608 (1984)).

Other modifications of the antibody are contemplated herein. For example, the antibody may be linked to one of a variety of nonproteinaceous polymers, *e.g.*, 35 polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol.

- 5 The antibodies disclosed herein may also be formulated as liposomes. Liposomes containing the antagonist are prepared by methods known in the art, such as described in Epstein *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:3688 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA*, 77:4030 (1980); U.S. Pat. Nos. 4,485,045 and 4,544,545; and W097/38731 published October 23, 1997.
- 10 Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

- Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG derivatized phosphatidylethanolamine (PEG-PE).
- 15 Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of an antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.* *J. Biol. Chem.* 257:286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See
- 20 Gabizon *et al.* *J. National Cancer Inst.* 81(19):1484 (1989).

- Amino acid sequence modification(s) of protein or peptide antagonists described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody are prepared by introducing appropriate
- 25 nucleotide changes into the antibody encoding nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antagonist. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses
- 30 the desired characteristics. The amino acid changes also may alter post-translational processes of the antagonist, such as changing the number or position of glycosylation sites.

- A useful method for identification of certain residues or regions of the antibody that are preferred locations for mutagenesis is called "alanine scanning
- 35 mutagenesis" as described by Cunningham and Wells *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (*e.g.*, charged

5 residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, 10 while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed antagonist variants are screened for the desired activity.

15 Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antagonist with an N-terminal methionyl residue or the antagonist fused to a 20 cytotoxic polypeptide. Other insertional variants of the antagonist molecule include the fusion to the N- or C-terminus of the antagonist of an enzyme, or a polypeptide which increases the serum half-life of the antagonist.

Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antagonist molecule 25 replaced by different residue. The sites of greatest interest for substitutional mutagenesis of antibody antagonists include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated 30 "exemplary substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Table 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gin; asn	lys
Asn (N)	gln; his; asp, lys; arg	gln
Asp (D)	glu; asn	glu
Cys (C)	ser; ala	ser
Gln (Q)	asn; glu	asn
Glu (E)	asp; gin	asp
Gly (G)	ala	ala
His (H)	asn; gin; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	tyr
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

- 5 Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side

5 chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophobic: cys, ser, thr;
- (3) acidic: asp, glu;
- 10 (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

15 Any cysteine residue not involved in maintaining the proper conformation of the antagonist also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bonds) may be added to the antagonist to improve its stability (particularly where the antagonist is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.* a humanized or human antibody). Generally, the resulting variants selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for
25 generating such substitutional variants is affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.* 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage
30 particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.* binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identified hypervariable region residues contributing
35 significantly to antigen binding. Alternatively, or in addition, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to

5 identify contact points between the antibody and antigen. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further
10 development.

Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antagonist. By altering is meant deleting one or more carbohydrate moieties found in the antagonist, and/or adding one or more
15 glycosylation sites that are not present in the antagonist.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the
20 recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most
25 commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation
30 sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antagonist (for O-linked glycosylation sites).

Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by a variety of methods known in the art. These methods
35 include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by

- 5 oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antagonist.

It may be desirable to modify the antibodies used in the invention to improve effector function, e.g. so as to enhance antigen-dependent
10 cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antagonist. This may be achieved by introducing one or more amino acid substitutions in an Fc region of an antibody antagonist. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric
15 antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shopes, *B. J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional
20 cross-linkers as described in Wolff *et al.* *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.* *Anti-Cancer Drug Design* 3:2 19-230 (1989).

To increase the serum half life of the antibody, one may incorporate a
25 salvage receptor binding epitope into the antagonist (especially an antibody fragment) as described in US Patent 5,739,277, for example. As used herein, the term "salvage
receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the
30 *in vivo* serum half-life of the IgG molecule.

IV. Pharmaceutical Formulations

Therapeutic formulations comprising antagonists used in accordance with the present invention are prepared for storage by mixing an antagonist
35 having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th

5 edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl
10 ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic
15 polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein
20 complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

The immunomodulatory antibody and the B cell depleting antibody may be in the same formulation or may be administered in different formulations. The composition may further include other non-antibody antagonists, e.g.,
25 CD40L or B7 antagonists. Examples thereof include soluble CD40, B7 and fusions thereof. Administration can be concurrent or sequential, and may be effective in either order.

Exemplary anti-CD20 antibody formulations are described in W098/56418, expressly incorporated herein by reference. This publication
30 describes a liquid multidose formulation comprising 40 mg/mL rituximab, 25 mM acetate, 150 mM trehalose, 0.9% benzyl alcohol, 0.02% polysorbate 20 at pH 5.0 that has a minimum shelf life of two years storage at 2-8°C. Another anti-CD20 formulation of interest comprises 10mg/mL rituximab in 9.0 mg/mL sodium chloride, 7.35 mg/mL sodium citrate dihydrate, 0.7mg/mL polysorbate
35 80, and Sterile Water for Injection, pH 6.5.

5 Lyophilized formulations adapted for subcutaneous administration are described in W097/04801. Such lyophilized formulations may be reconstituted with a suitable diluent to a high protein concentration and the reconstituted formulation may be administered subcutaneously to the mammal to be treated herein.

10 The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide a chemotherapeutic agent, cytokine or immunosuppressive agent (*e.g.* one which acts on T cells, such as
15 cyclosporin or an antibody that binds T cells, *e.g.* one which binds LFA-1). The effective amount of such other agents depends on the amount of antagonist present in the formulation, the type of disease or disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the
20 heretofore employed dosages.

 The active ingredients may also be entrapped in microcapsules prepared, for example, by 30 coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate)
25 microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

 Sustained-release preparations may be prepared. Suitable examples of
30 sustained release preparations include semipermeable matrices of solid hydrophobic polymers containing the antagonist, which matrices are in the form of shaped articles, *e.g.* films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S.
35 Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid

5 copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

10

V. Treatment with the B Cell Depleting Antibody and Immunoregulatory Antibody

A composition comprising B cell depleting antibody and/or an immunoregulatory antibody will be formulated, dosed, and administered in a
15 fashion consistent with good medical practice. Factors for consideration in this context include the particular B cell malignancy or disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disease or disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to
20 medical practitioners. The therapeutically effective amount of the antagonist to be administered will be governed by such considerations.

As noted previously, the B cell depleting antibody and the immunoregulatory antibody may be in the same or in different formulations. These antagonist formulations can be administered separately or concurrently,
25 and in either order. Preferably, the B cell depleting antibody specific to the B cell antigen target, *e.g.*, CD20, CD19, CD22, CD37 or CD22, will be administered separately from the immunoregulatory antibody, *e.g.*, an anti-CD40L antibody, anti-CD40 antibody, or anti-B7 antibody. Preferably, the CD40L antibody will be the humanized anti-CD40L antibody disclosed in U.S.
30 Patent 6,001,358 and the anti-B7 antibody the primatized antibody disclosed in US Patent 6,113,898. As noted, this antibody has recently been shown to possess apoptotic activity. Also the preferred CD40L antibody has been shown to have efficacy in treatment of both T and B cell autoimmune diseases. Also, unlike another humanized anti-CD40L antibody (5c8) reported by Biogen, this
35 antibody is not known to cause any adverse toxicity.

5 As a general proposition, the therapeutically effective amount of an antibody administered parenterally per dose will typically be in the range of about 0.1 to 500 mg/kg of patient body weight per day, with the typical initial range of antagonist used being in the range of about 2 to 100 mg/kg.

 The preferred B cell depleting antibody is RITUXAN®. Suitable
10 dosages for such antibody are, for example, in the range from about 20mg/m² to about 1000mg/m². The dosage of the antibody may be the same or different from that presently recommended for RITUXAN® for the treatment of non-Hodgkin's lymphoma. For example, one may administer to the patient one or more doses of substantially less than 375mg/m² of the antibody, *e.g.* where the
15 dose is in the range from about 20mg/m² to about 250mg/m², for example from about 50mg/m² to about 200mg/m².

 Moreover, one may administer one or more initial doses) of the antibody followed by one or more subsequent dose(s), wherein the mg/m² dose of the antibody in the subsequent doses) exceeds the mg/m² dose of the antibody in the
20 initial dose(s). For example, the initial dose may be in the range from about 20mg/m² to about 250mg/m² (*e.g.* from about 50mg/m² to about 200mg/m²) and the subsequent dose may be in the range from about 250mg/m² to about 1000mg/m².

 As noted above, however, these suggested amounts of both
25 immunoregulatory and B cell depleting antibody are subject to a great deal of therapeutic discretion. The key factor in selecting an appropriate dose and scheduling is the result obtained, as indicated above. For example, relatively higher doses may be needed initially for the treatment of ongoing and acute diseases. To obtain the most efficacious results, depending on the particular B
30 cell malignancy, the antagonist is administered as close to the first sign, diagnosis, appearance, or occurrence of the disease or disorder as possible or during remissions of the disease or disorder.

 The antibodies are administered by any suitable means, including parenteral,
35 subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local

5 immunosuppressive treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the antibody may suitably be administered by pulse infusion, *e.g.*, with declining doses of the antibody. Preferably the dosing is given by injections, most preferably intravenous or
10 subcutaneous injections, depending in part on whether the administration is brief or chronic.

One additionally may administer other compounds, such as chemotherapeutic agents, immunosuppressive agents and/or cytokines with the antibodies herein. The combined administration includes co-administration,
15 using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

Aside from administration of antibodies to the patient the present
20 application contemplates administration of antibodies by gene therapy. Such administration of nucleic acid encoding the antibodies is encompassed by the expression "administering a therapeutically effective amount of an antagonist". See, for example, W096/07321 published March 14, 1996 concerning the use of gene therapy to generate intracellular antibodies.

25 There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells; *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the site where the antagonist is required. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells and the
30 modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes which are implanted into the patient (see, *e.g.* U.S. Patent Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into
35 cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the

5 use of liposomes, electroporation, microinjection, cell fusion, DEAF-dextran, the calcium phosphate precipitation method, etc. A commonly used vector for ex vivo delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or
10 adeno associated virus) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are DOTMA, DOPE and DC-Chol, for example). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane
15 protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.* capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, and
20 proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, *J. Biol. Chem* 262:4429-4432 (1987); and Wagner *et al.*, *Proc. Natl. Acad. Sci. USA* 87:3410-3414(1990). For review of the currently known gene marking and gene therapy protocols see Anderson *et al.*, *Science* 256:808-8 13
25 (1992). See also WO 93/25673 and the references cited therein.

-VI. Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the diseases or disorders
30 described above is provided.

The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds or contains a
35 composition which is effective for treating the disease or disorder of choice and may have a sterile access port (for example the container may be an intravenous

5 solution bag or a vial having a stopper pierceable by a hypodermic injection
needle). As whole, there may be one or several compositions. At least one
active agent in one of those compositions is an antibody having B cell depleting
activity and at least one antibody is an immunoregulatory antibody such as an
anti-CD40L, anti-CD40, anti-CD23, anti-CD4 or anti-B7 antibody. The label or
10 package insert indicates that the composition is used for treating a patient
having or predisposed to B cell malignancy, such as those listed hereinabove.
The article of manufacture may further comprise a second container comprising
a pharmaceutically acceptable buffer, such as bacteriostatic water for injection
(BWF), phosphate-buffered saline, Ringer's solution and dextrose solution. It
15 may further include other materials desirable from a commercial and user
standpoint, including other buffers, diluents, filters, needles, and syringes.

Further details of the invention are illustrated by the following
non-limiting Examples. The disclosures of all citations in the specification are
expressly incorporated herein by reference.

20 The antibodies of the invention may be administered to a human or other
animal in accordance with the aforementioned methods of treatment in an
amount sufficient to produce such effect to a therapeutic or prophylactic degree.
Such antibodies of the invention can be administered to such human or other
animal in a conventional dosage form prepared by combining the antibody of
25 the invention with a conventional pharmaceutically acceptable carrier or diluent
according to known techniques. It will be recognized by one of skill in the art
that the form and character of the pharmaceutically acceptable carrier or diluent
is dictated by the amount of active ingredient with which it is to be combined,
the route of administration and other well-known variables.

30 The routine of administration of the antibody (or fragment thereof) of
the invention may be oral, parenteral, by inhalation or topical. The term
parenteral as used herein includes intravenous, intraperitoneal, intramuscular,
subcutaneous, rectal or vaginal administration. The subcutaneous and
intramuscular forms of parenteral administration are generally preferred.

35 The daily parenteral and oral dosage regimes for employing compounds
of the invention to prophylactically or therapeutically induce

- 5 immunosuppression, or to therapeutically treat carcinogenic tumors will generally be in the range of about 0.05 to 100, but preferably about 0.5 to 10, milligrams per kilogram body weight per day.

The antibodies of the invention may also be administered by inhalation. By "inhalation" is meant intranasal and oral inhalation administration.

- 10 Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The preferred dosage amount of a compound of the invention to be employed is generally within the range of about 10 to 100 milligrams.

- The antibodies of the invention may also be administered topically. By
15 topical administration is meant non-systemic administration and includes the application of an antibody (or fragment thereof) compound of the invention externally to the epidermis, to the buccal cavity and instillation of such an antibody into the ear, eye and nose, and where it does not significantly enter the blood stream. By systemic administration is meant oral, intravenous,
20 intraperitoneal and intramuscular administration. The amount of an antibody required for therapeutic or prophylactic effect will, of course, vary with the antibody chosen, the nature and severity of the condition being treated and the animal.

25 EXAMPLES

Example 1

Properties of B lymphoma cells. DHL-4 cells

- The concept that anti-CD40L antibody could block CD40L-CD40 mediated survival of malignant B-cells from chemotherapy induced
30 toxicity/apoptosis was tested *in vitro* using IDEC-131, and the B-lymphoma cell line, DHL-4 (Roos et al., *Leuk. Res.* 10: 195-202 (1986)) exposed to adriamycin (ADM). IDEC-131 is a humanized version of the murine, monoclonal anti-human CD40L antibody, 24-31.

- Initially, the minimum concentration of ADM cytotoxic to DHL-4 cells
35 was determined by exposing DHL-4 cells for 4 hours to different concentrations of ADM. The cell cytotoxicity of DHL-4 cells after 5 days in culture was

5 measured by Alamar Blue, a dye-reduction assay by live cells (see
Gazzano-Santoro *et al.*, *J. Immunol. Meth.* 202: 163-171 (1997)). Briefly, 1×10^5 DHL-4 cells in growth medium (RMPI-1640 plus 10% Fetal Calf Serum)
were incubated with varying concentrations of ADM (1×10^{-6} M to 1×10^{-8} M)
10 in cell culture tubes at 37 °C. for 4 hours. After incubation, cells were washed,
re-suspended in growth medium at 1×10^5 cells/ml concentration and 200 μ l of
cell suspension was added to each well of 96-well flat-bottom plate. Plates were
incubated at 37 °C. and tested for cytotoxicity at different time points. During
the last 18 hours of incubation, 50 μ l of redox dye Alamar Blue (Biosource
15 International, Cat. #DAL 1100) was added to each well. Following incubation,
plates were cooled by incubating at room temperature for 10 minutes on a
shaker, and the intracellular reduction of the dye was determined. Fluorescence
was read using a 96-well fluorometer with excitation at 530 nm and emission at
590 nm. The results are expressed as relative fluorescence units (RFU). The
percentage of cytotoxicity was calculated as follows:

20
$$[1 - (\text{average RFU of test sample} \div \text{Average RFU of control cells})] \times 100\%.$$
 Titration curve of ADM cytotoxicity was established and minimal
concentrations of the drug for cytotoxicity was selected for subsequent assays.

The results, as displayed in Fig. 1, shows cell cytotoxicity of DHL-4
cells cultured for 5 days after being exposed to ADM (2×10^{-7} M and 4×10^{-8} M
25 of ADM) for 4 hours prior to culture. Cells were washed once after exposure
and cultured in growth medium for 5 days and cytotoxicity determined by
Alamar Blue dye-uptake assay, as described above. Additionally, the DHL-4
cells were characterized for the membrane expression of selected CD molecules
by flow cytometry. DHL-4 cells have been found to express CD19, CD20,
30 CD40 molecules, but no expression of CD40L was detected.

Example 2

Anti -CD40L antibody overrides CD40L mediated resistance to killing by to killing, by adriamycin in of -lymphoma cells

35 Fig. 2A shows the effect of an anti-CD40L antibody on CD40L-CD40
mediated resistance of DHL-4 cells to cell death induced by ADM. DHL-4

5 cells (0.5×10^6 cells/ml) were incubated in the presence of 10 $\mu\text{g/ml}$ of soluble CD40L (sCD40L, P. A. Brams, E. A. Padlan, K. Hariharan, K. Slater, J. Leonard, R. Noelle, and R. Newman, "A humanized anti-human CD 154 monoclonal antibody blocks CD 154-CD40 mediated human B cell activation," (*manuscript submitted*)) for 1 hour at 37°C. After 1 hour of incubation, low
10 concentrations of ADM (2×10^{-7} M - 4×10^{-8} M) were added and incubated for another 4 hours in the presence or absence of CD40L (10 $\mu\text{g/ml}$). Following exposure to ADM, cells were washed and resuspended in growth medium at 0.5×10^6 cells/ml concentration, and 100 μl of cell suspension added to each well of
15 96-well flat bottom plate, in duplicate, with or without sCD40L. sCD40L (10 $\mu\text{g/ml}$) was added to cultures that have been continuously exposed to sCD40L during ADM treatment and to cultures that had no sCD40L during ADM exposure. In addition, IDEC-131 at 10 $\mu\text{g/ml}$ was added to cultures to determine its effect on DHL-4 cells incubated with sCD40L and ADM. After 5 days, the cytotoxicity was measured by Alamar Blue dye-uptake assay, as
20 described.

Data show that sCD40L prolonged survival of DHL-4 cells after ADM treatment, whereas, as expected, increased cytotoxicity was observed in cells that were exposed to ADM in the absence of sCD40L. Furthermore, addition of anti-CD40L antibody (IDEC-131) reversed CD40L mediated cell survival,
25 leading to increase in cell cytotoxicity (Fig. 2A).

The addition of IDEC-131 alone had no effect on DHL-4 cells treated with sCD40L, which indicates that the antibody, by itself, does not have any direct inhibitory or cytotoxic activities on DHL-4 cells (Fig. 2B). DHL-4 cells pre-incubated with and without sCD40L were cultured in the presence of
30 different concentrations of IDEC-131, RITUXAN®, the anti-CD20 antibody CE9.1, and anti-CD4 antibodies (Anderson et al., *Clin. Immunol. & Immunopathol.* 84: 73-84 (1997)). After 5 days, the cytotoxicity/proliferation of DHL-4 cells was determined by Alamar Blue assay, as described above. Fig. 2B shows no effect on the proliferation or the cytotoxicity of DHL-4 cells by
35 IDEC-131, whereas RITUXAN®, as expected, inhibited cell proliferation and

- 5 induced cytotoxicity. No effect was seen in the DHL-4 cells cultured with anti-CD4 antibodies.

Example 3

CD40L-CD40 signaling prevents apoptosis of B-lymphoma
10 cells by anti-CD20 antibody, RITUXAN®

The effect of CD40L-CD40 mediated signaling on anti-CD20 antibody induced apoptosis of B-lymphoma cells was determined using an *in vitro* system involving DHL-4 cells and the surface cross-linking of RITUXAN®. DHL-4 cells (0.5 to 1×10^6 cells/ml) were cultured with sCD40L ($10 \mu\text{g/ml}$) at 15 37°C . After overnight culture, cells were harvested and incubated with $10 \mu\text{g/ml}$ of RITUXAN® or the control antibody (CE9.1; an anti-CD4 antibody) with or without sCD40L ($10 \mu\text{g/ml}$) on ice. After 1 hour of incubation, cells were centrifuged to remove unbound antibodies, and resuspended at 1×10^6 cells/ml in growth medium (5% FCS-RPMI) and cultured in tissue culture 20 tubes. The cells surface bound antibodies were cross-linked by spiking F(ab')_2 fragments of goat anti-human Ig-Fc γ specific antibodies at $15 \mu\text{g/ml}$, and the cultures were incubated at 37°C . until assayed for apoptosis. Apoptosis was detected using a flow cytometry caspase-3 assay. Cultured cells were harvested at 4 and 24 hours, washed and fixed at 4°C . using Cytofix (Cytofix/Cytoperm™ 25 Kit, Pharmingen Cat. #2075KK). After 20 min of fixation, cells were washed and $15 \mu\text{l}$ of affinity purified PE-conjugated polyclonal rabbit anti-caspase-3 antibody (Pharmingen, Cat. # 67345) and $50 \mu\text{l}$ of cytoperm (Pharmingen; Cat. #2075KK) were added. Cells were incubated on ice in the dark for 30 min. After incubation cells were washed once and resuspended in cytoperm. Flow 30 cytometry data was acquired on FACScan and analyzed using WinList software from Verity Software House.

Table I shows resistance of RITUXAN® induced apoptosis in DHL-4 lymphoma cells by exposure to sCD40L. In these studies, activation of caspase-3 was used as the surrogate marker since our previous studies revealed 35 good correlation between caspase-3 and Tunel assay. Cross-linking of RITUXAN® on the DHL-4 cell surface in the presence of sCD40L decreased

- 5 levels of apoptosis, whereas cells not exposed to sCD40L apoptosed. In comparison, cultures incubated in the presence of an antibody of the same isotype, control antibody (CE9.1), resulted in no apoptosis of the cells. Thus, the data suggests that sCD40L induced signaling of CD40 pathway can lead to development of RITUXAN® mediated killing of B-lymphoma cells.

10

Table I:

Resistance of RITUXAN® mediated apoptosis of DHL-4 cells by sCD40L

Culture Conditions	% Apoptosis (IVHF) ^(a)	
	4 Hours	24 Hours
<u>DHT-4 cells exposed to sCD40L</u>		
Cells only	3.35 (17.42)	4.94 (7.62)
Cells + RITUXAN	1.97 (1.97)	4.54 (6.54)
Cells + RITUXAN + anti-hu.IgG.F(ab') ₂	21.17 (17.39)	9.62 (13.44)
Cells + CE9.1	2.31 (13.25)	4.15 (7.85)
Cells + CE9.1 + anti-hu.IgG.F(ab') ₂	2.09 (22.14)	4.14 (9.57)
Cells + anti-hu.IgG.F(ab') ₂	1.93 (12.57)	5.13 (8.02)
<u>DHL-4 cells not exposed to sCD40L</u>		
Cells only	4.36 (14.34)	5.08 (17.62)
Cells + RITUXAN	5.67 (10.66)	1.08 (17.92)
Cells + RITUXAN + anti-hu.IgG.F(ab') ₂	74.82 (22.80)	30.63 (26.84)
Cells + CE9.1	5.99 (14.00)	3.05 (18.24)
Cells + CE9.1 + anti-hu.I-G.F(ab') ₂	5.96 (12.11)	2.24 (18.19)
Cells + anti-hu.IgG.F(ab') ₂	6.09 (12.27)	1.85 (17.27)

^(a) Percent positive cells with caspase-3 activity and its mean fluorescent intensity in log scale.

15

Example 4**Effect of IDEC-131 on the survival of chronic lymphocytic leukemia (CLL) cells**

- To determine the effect of IDEC-131 on the growth and survival of B-CLL cells *in vitro*, B-CLL cells were cultured with and without IDEC-131 in the presence of CD40L *in vitro*. Peripheral blood mononuclear cells (PBMC) were isolated from a CLL patient's blood using a Ficoll-Hypaque gradient centrifugation. Viability was determined by Trypan blue dye exclusion and was >98%. Flow cytometric analysis revealed that >70% of the lymphocytes were CD 19⁺/CD20⁺. CLL cells (PBMC) were cultured in CLL growth medium

- 5 (e.g., RPMI-1640 medium supplemented with 5% FCS or 2% of autologous donor plasma, supplemented with 2 mM L-Glutamine and 100 U/ml Penicillin-Streptomycin). In addition, for some experiments, CD19⁺ B-cells were purified using CD19⁺ Dynabeads™ as per manufacture's instructions (Dynal, Cat. #111.03/111.04) and cultured as above. CLL or purified B-CLL
- 10 cells cultured in growth medium mostly under went spontaneous apoptotic cell death. However, culturing these cells in the presence of sCD40L extended their viability in cultures. Table II indicates the cell viability of CD 19⁺ B-CLL cells grown in the presence or absence of sCD40L (5 µg/ml) at different time points and indicates the longer survival of CLL cells. B-CLL cells from Patient #1
- 15 cultured with sCD40L had ≥ 60% viability for greater than 2 weeks, whereas cells grown in the absence of sCD40L had less than 10% viability.

Table II:

Survival of B-CLL cells in the presence of sCD40L

B-CLL Sample	Time (Hours)	%Viability ^(a)	
		(-) CD40L	(+) CD40L
Patient #1	0	≥90	≥90
	48	88	90
	96	46	77
	144	30	72
Patient #2	0	≥90	≥90
	72	40	72
	96	31	65
	144	17	51

^(a) equals the percent viability determined by Trypan blue dye exclusion.

- 20 Fig. 3A shows the effect of IDEC-131 on the growth and survival of B-CLL cells after 7 days in culture. Purified B-CLL cells from a CLL patient (2 x 10⁶ cells/ml) were divided into two culture tubes. Cells in one tube were mixed with sCD40L (5 µg/ml) in equal volume of growth medium, whereas the other tube was incubated with equal volume of growth medium as control. After 1
- 25 hour of incubation at 37° C., cells were gently mixed and 100 µl of cell suspension media added to each well of a 96-well flat bottom plate in duplicate with and without varying concentrations of IDEC-131 (10 µg/ml to 0.3 µg/ml). Seven days later, cell survival/death in culture was determined by Alamar Blue assay, as described above. Data showed cell survival in cultures with sCD40L.

- 5 The addition of IDEC-131 into culture resulted in increased cell death, which indicated a reversal of cell survival or a sensitization to cell death. Additionally, RITUXAN® administered at the same concentration as the IDEC-131 produced less of lower effect than IDEC-131 on cell death (Fig. 3B).

10

Example 5

CD40L-CD40 mediated up-regulation of HLA-DR molecules in B-CLL

- To determine whether the CD40L-CD40 signal transduction pathway is intact, CLL cells from CLL patients were cultured (5×10^5 cells/ml) with and without $5 \mu\text{g/ml}$ of CD40L at 37°C . At 48 hours and 144 hours, the class II molecule, HLA-DR expression, was determined on $\text{CD } 19^+$ cells by flow cytometry using standard procedures. Briefly, cultured lymphocytes were harvested at different time points and analyzed for surface expression of molecules using antibodies coupled to either fluorescein (FITC) or phycoerythrin (PE) for single or double staining using a FACScan (Becton-Dickinson) flow cytometer. To stain for flow cytometry, 1×10^6 cells in culture tubes were incubated with appropriate antibodies as follows: anti-CD45-FITC to gate lymphocyte population on a scatter plot; anti-CD19-PE (Pharmingen, Cat. # 30655) or anti-CD20-FITC (Pharmingen; Cat. #33264) antibodies to determine the $\text{CD } 19^+$ and/or $\text{CD } 20^+$ B-cells; anti-CD3-FITC antibodies (Pharmingen; Cat. #30104) to gate-off the T cells; anti-CD 19-RPE and anti-HLA-DR-FITC antibodies (Pharmingen; Cat. #32384) to determine the Pclass II expression on $\text{CD } 19^+$ cells. Cells were washed once by centrifugation (at $200 \times g$, for 6 min.) with 2 ml cold PBS and incubated with antibody for 30 min. on ice, after which the cells washed once, fixed in 0.5% paraformaldehyde and stored at 4°C . until analyzed. Flow cytometry data was acquired on FACScan and analyzed using WinList software (Verity Software House). The machine was set to autogating to allow examination of quadrants containing cells that were single stained with either RPE or FITC, unstained or doubly stained. Fig. 4 shows the comparison of HLA-DR expression in $\text{CD } 19^+$ CLL cells cultured with sCD40L and those cells not cultured with sCD40L. A higher

- 5 level of HLA-DR expression was detected on B-CLL cells cultured in the presence of sCD40L (Table III).

Table III

CD40L-CD40 mediated up-regulation of HLA-DR molecule in B-CLL

Sample	Time	HLA-DR ^{+(a)}	
		%Positive	MFI
Control	48 hrs	81	92
	144 hrs	88	1655
Cells + sCD40L	48 hrs	88	101
	144 hrs	95	2943

- (a) CD19⁺ B-cells that are positive for HLA-DR molecules and its mean
10 fluorescent intensity (MFI).
(b)

Example 6

Preparation of IDEC-131 and RITUXAN®

For treatment of a CD40⁺ malignancy, IDEC-131 at about 10 to about
15 50 mg/ml in a formulation buffer 10 mM Na-citrate, 150 mM NaCl, 0.02% Polysorbate 80 at pH 6.5 is infused intravenously (iv) to a subject. IDEC-131 is administered before, after or in conjunction with RITUXAN®. The RITUXAN® dosage infused ranges from about 3 to about 10 mg/kg of subject weight.

20

Example 7

Preparation of IDEC-131 and CHOP

For treatment of CD40⁺ malignancies responsive to CHOP (e.g., Hodgkin's Disease, Non-Hodgkin's lymphoma and chronic lymphocytic
25 leukemia, as well as salvage therapy for malignancies wherein cells are CD40⁺), IDEC-131 is infused at a dosage ranging from about 3 to about 10 mg per kg of patient weight immediately prior to the initiation of the CHOP cycle. IDEC-131 administration will be repeated prior to each CHOP cycle for a total of 4 to 8 cycles.

30

5

Example 8Administration of anti- CD40L or anti-B7 in combination with RITUXAN®
to treat B-cell lymphoma in a subject

Combination therapies are particularly useful as salvage therapies or for treating relapsed or aggressive forms of CD40⁺ malignancies (e.g., Hodgkin's Disease, Non-Hodgkin's lymphoma and CLL). When IDEC-131 is to be administered in combination with CHOP and RITUXAN®, IDEC-131 is administered as discussed above in Example 6, followed by the schedule specified for CHOP-IDEC-131 administration in Example 7. Alternatively, the same regimen is effected wherein IDEC-131 (anti-CD40L) is substantially within an anti-B7 antibody.

All references discussed above are hereby incorporated by reference in their entirety.

5

CLAIMS

1. A method for treating CD40⁺ malignancies comprising administering a therapeutically effective amount of an antibody or antibody fragment which binds to CD40L thereby inhibiting CD40/CD40L interaction or CD40 signaling.
2. The method of Claim 1, wherein the CD40⁺ malignancy is a B-cell lymphoma or a B-cell leukemia.
3. The method of Claim 2, wherein the B-cell lymphoma is Hodgkin's Disease (HD) or Non-Hodgkin's Lymphoma (NHL).
4. The method of Claim 3, wherein the NHL is low grade, intermediate grade or high grade.
5. The method of Claim 3, wherein the NHL is selected from the subtype group consisting of: small lymphocytic, follicular and predominantly small cleaved cell, follicular and mixed small cleaved and large cell type, follicular and predominantly large cell type, diffuse small cleaved cell, diffuse mixed small and large cell, diffuse large cell, large cell immunoblastic, lymphoblastic, small non-cleaved Burkitt's and non-Burkitt's type, AIDS-related lymphomas, angioimmunoblastic lymphadenopathy, mantle cell lymphoma, and monocytoid B-cell lymphoma.
6. The method of Claim 2, wherein the B-cell leukemia is a chronic B-cell leukemia, acute lymphoblastic leukemia of a B-cell lineage, or chronic lymphocytic leukemia of a B-cell lineage.
7. The method of Claim 2, wherein the antibody or antibody fragment which binds to CD40L is IDEC-131, 3E4, 2H5, 2H8, 4D9-8, 4D9-9, 24-31, 24-43, 89-76 or 89-79.

5 8. The method of Claim 7, wherein the antibody or antibody
fragment is chimeric, bispecific, human or humanized.

 9. The method of Claim 2, wherein the antibody fragment is Fab,
Fab', scFv or F(ab')₂.

10

 10. The method of Claim 2, further comprising administering a
therapeutically effective amount of a second antibody or fragment thereof, a
chemotherapeutic, a combination of chemotherapeutic agents and/or a
radiotherapy.

15

 11. The method of Claim 10, wherein the radiotherapy is external
radiation treatment or a radiolabeled antibody.

 12. The method of Claim 11, wherein the radiolabeled antibody is
20 radiolabeled IDEC-131, RITUXAN®, or B1 or fragments thereof.

 13. The method of Claim 12, wherein the radiolabeled antibody is
radiolabeled with ¹²³I, ¹²⁵I, ¹³¹I, ¹¹¹In, ¹³¹In, ³²P, ⁶⁴Cu, ⁶⁷Cu, ²¹¹At, ¹⁷⁷Lu, ⁹⁰Y,
¹⁸⁶Re, ²¹²Pb, ²¹²Bi, ⁴⁷Sc, ¹⁰⁵Rh, ¹⁰⁹Pd, ¹⁵³Sm, ¹⁸⁸Re, ¹⁹⁹Au, ²¹¹At, and ²¹³Bi.

25

 14. The method of Claim 10, wherein the chemotherapeutic agent for
treating HD is any one or more of the following: an alkylating agent, a vinca
alkaloid, procarbazine, methotrexate or prednisone.

30 15. The method of Claim 10, wherein the chemotherapeutic agent for
treating NHL is any one or more of the following: an alkylating agent,
cyclophosphamide, chlorambucil, 2-CDA, 2'-deoxycoformycin, fludarabine,
cytosine arabinoside, cisplatin, etoposide or ifosfamide.

35 16. The method of Claim 10, wherein the combination of
chemotherapeutic agents for treating HD is: MOPP, ABVD, CH₁VPP, CABS,

- 5 MOPP plus ABVD, MOPP plus ABV, BCVPP, VABCD, ABDIC, CBVD, PCVP, CEP, EVA, MOPLACE, MIME, MINE, CEM, MTX-CHOP, EVAP or EPOCH.

- 10 17. The method of Claim 10, wherein the combination of chemotherapeutic agents for treating NHL is: CVP, CHOP, C-MOPP, CAP-BOP, m-BACOD, ProMACE-MOPP, ProMACE-CytaBOM, MACOP-B, IMVP-16, MIME, DHAP, ESHAP, CEPP(B) or CAMP.

- 15 18. The method of Claim 10, wherein the chemotherapeutic agent for treating a B-cell leukemia is at least one of the following: anthracycline, cyclophosphamide, L-asparaginase and a purine analog.

- 20 19. The method of Claim 10, wherein the combination of chemotherapeutic agents for treating a B-cell leukemia is: vincristine, prednisone, anthracycline and cyclophosphamide or asparaginase; vincristine, prednisone, anthracycline, cyclophosphamide and asparaginase; CHOP; CMP; CVP; COP or CAP.

- 25 20. The method of Claim 10, wherein the second antibody is selected from the group consisting of an anti-CD20 antibody, anti-CD19 antibody, anti-CD22 antibody, and anti-CD40 antibody.

- 30 21. The method of Claim 21, wherein the anti-CD20 antibody is RITUXAN® or a fragment thereof or B1 or a fragment thereof.

- 35 22. A method of treating a CD40⁺ malignancy comprising the step of administering an anti-CD40L antibody or fragment thereof wherein the anti-CD40L antibody or antibody fragment blocks CD40-CD40L interaction or inhibits CD40 signaling; and administering a second antibody or fragment selected from the group consisting of an anti-CD20, anti-CD40, anti-CD19, and anti-CD22 antibody or fragment thereof.

5 23. The method of Claim 22, wherein the CD40⁺ malignancy is a B-cell lymphoma or a B-cell leukemia.

 24. A combination therapy for the treatment of a CD40⁺ malignancy comprising a CD40L antagonist and at least one of the following (a) a
10 chemotherapeutic agent or a combination of chemotherapeutic agents, (b) a radiotherapy, (c) an anti-CD20 antibody or fragment thereof and (d) anti-CD40 antibody or fragment thereof, (e) an anti-CD19 antibody or fragment thereof, and (f) an anti-CD22 antibody or fragment thereof.

15 25. The method of Claim 24, wherein the radiotherapy is external radiation treatment or a radiolabeled antibody.

 26. The method of Claim 25, wherein the radiolabeled antibody is radiolabeled with ¹²³I, ¹²⁵I, ¹³¹I, ¹¹¹In, ¹³¹In, ³²P, ⁶⁴Cu, ⁶⁷Cu, ²¹¹At, ¹⁷⁷Lu, ⁹⁰Y, ¹⁸⁶Re, ²¹²Pb, ²¹²Bi, ⁴⁷Sc, ¹⁰⁵Rh, ¹⁰⁹Pd, ¹⁵³Sm, ¹⁸⁸Re, ¹⁹⁹Au, ²¹¹At, and ²¹³Bi.
20

 27. The combination therapy of Claim 24 wherein the CD40⁺ malignancy is a B-cell leukemia or B-cell lymphoma.

25 28. The combination therapy of Claim 27, wherein the B-cell lymphoma is HD or NHL.

 29. The combination therapy of Claim 28, wherein the NHL is low grade, intermediate grade or high grade.

30

 30. The combination therapy of Claim 28, wherein the NHL is selected from the subtype group consisting of the following: small lymphocytic, follicular and predominantly small cleaved cell, follicular and mixed small cleaved and large cell type, follicular and predominantly large cell type, diffuse
35 small cleaved cell, diffuse mixed small and large cell, diffuse large cell, large

5 cell immunoblastic, lymphoblastic, small non-cleaved Burkitt's and non-Burkitt's type, AIDS-related lymphomas, angioimmunoblastic lymphadenopathy, mantle cell lymphoma and monocytoid B-cell lymphoma.

31. The combination therapy of Claim 28, wherein the B-cell
10 leukemia is a chronic B-cell leukemia, acute lymphoblastic leukemia of a B-cell lineage, or chronic lymphocytic leukemia of a B-cell lineage.

32. The combination therapy of Claim 24, wherein the CD40L
antagonist is an anti-CD40L antibody or a fragment thereof.

15 33. The combination therapy of Claim 32, wherein the anti-CD40L antibody is IDEC-131 or a fragment thereof.

34. The combination therapy of Claim 32, wherein the anti-CD40L
20 fragment is Fab, Fab', scFv or F(ab')₂.

35. The combination therapy of Claim 24, wherein the anti-CD20 antibody is RITUXAN® or a fragment thereof or B1 or a fragment thereof.

25 36. The combination therapy of Claim 28, wherein the chemotherapeutic agent for treating HD is any one or more of the following: an alkylating agent, a vinca alkaloid, procarbazine, methotrexate or prednisone.

37. The combination therapy of Claim 28, wherein the
30 chemotherapeutic agent for treating NHL is any one or more of the following: an alkylating agent, cyclophosphamide, chlorambucil, 2-CDA, 2'-deoxycoformycin, fludarabine, cytosine arabinoside, cisplatin, etoposide or ifosfamide.

35 38. The combination therapy of Claim 28, wherein the combination of chemotherapeutic agents for treating HD is: MOPP, ABVD, ChlVPP, CABS,

5 MOPP plus ABVD, MOPP plus ABV, BCVPP, VABCD, ABDIC, CBVD,
PCVP, CEP, EVA, MOPLACE, MIME, MINE, CEM, MTX-CHOP, EVAP or
EPOCH.

39. The combination therapy of Claim 28, wherein the combination
10 of chemotherapeutic agents for treating NHL is: CVP, CHOP, C-MOPP, CAP-
BOP, m-BACOD, ProMACE-MOPP, ProMACE-CytaBOM, MACOP-B,
IMVP-16, MIME, DHAP, ESHAP, CEPP(B), or CAMP.

40. The combination therapy of Claim 28, wherein the
15 chemotherapeutic agent for treating a B-cell leukemia is: anthracycline,
cyclophosphamide, L-asparaginase, a purine analog.

41. The combination therapy of Claim 28, wherein the combination
of chemotherapeutic agents for treating a B-cell leukemia is: vincristine,
20 prednisone, anthracycline and cyclophosphamide or asparaginase; vincristine,
prednisone, anthracycline, cyclophosphamide and asparaginase; CHOP; CMP;
CVP; COP or CAP.

42. A composition for the treatment of a CD40⁺ malignancy
25 comprising an (i) anti-CD40L antibody or antibody fragment thereof and at
least one of the following: (ii) a radiolabeled antibody that binds CD40L, CD19,
CD22, or CD20, (iii) an anti-CD20, an anti-CD19 antibody, an anti-CD22
antibody, or fragment thereof, or (iv) a chemotherapeutic agent or a
chemotherapeutic combination.

30

43. The composition for the treatment of a CD40⁺ malignancy of
Claim 42 wherein the malignancy is a B-cell lymphoma or a B-cell leukemia.

44. The composition of Claim 43, wherein the B-cell leukemia is
35 Hodgkin's Disease or NHL.

5 45. The composition of Claim 42, wherein the radiolabeled antibody is radiolabeled IDEC-131, RITUXAN®, or B1.

 46. The composition of Claim 46, wherein the radiolabeled antibody is radiolabeled with ¹²³I, ¹²⁵I, ¹³¹I, ¹¹¹In, ¹³¹In, ³²P, ⁶⁴Cu, ⁶⁷Cu, ²¹¹At, ¹⁷⁷Lu, ⁹⁰Y,
10 ¹⁸⁶Re, ²¹²Pb, ²¹²Bi, ⁴⁷Sc, ¹⁰⁵Rh, ¹⁰⁹Pd, ¹⁵³Sm, ¹⁸⁸Re, ¹⁹⁹Au, ²¹¹At, and ²¹³Bi.

 47. The composition of Claim 44, wherein the NHL is low grade, intermediate grade or high grade.

15 48. The composition of Claim 44, wherein the NHL is selected from the NHL subtype group consisting of the following: small lymphocytic, follicular and predominantly small cleaved cell, follicular and mixed small cleaved and large cell type, follicular and predominantly large cell type, diffuse small cleaved cell, diffuse mixed small and large cell, diffuse large cell, large
20 cell immunoblastic, lymphoblastic, small non-cleaved Burkitt's and non-Burkitt's type, AIDS-related lymphomas, angioimmunoblastic lymphadenopathy, mantle cell lymphoma and monocytoid B-cell lymphoma.

 49. The composition of Claim 42, wherein the anti-CD40L antibody
25 is IDEC-131 or a fragment thereof.

 50. The composition of Claim 42, wherein the anti-CD20 antibody is RITUXAN® or a fragment thereof or B1 or a fragment thereof.

30 51. The composition of Claim 43, wherein the chemotherapeutic agent for treating HD is any one or more of the following: an alkylating agent, a vinca alkaloid, procarbazine, methotrexate or prednisone.

 52. The composition of Claim 44, wherein the chemotherapeutic
35 agent for treating NHL is any one or more of the following: an alkylating agent,

5 cyclophosphamide, chlorambucil, 2-CDA, 2'-deoxycoformycin, fludarabine, cytosine arabinoside, cisplatin, etoposide or ifosfamide.

53. The composition of Claim 44, wherein the combination of chemotherapeutic agents for treating HD is: MOPP, ABVD, ChIVPP, CABS,
10 MOPP plus ABVD, MOPP plus ABV, BCVPP, VABCD, ABDIC, CBVD, PCVP, CEP, EVA, MOPLACE, MIME, MINE, CEM, MTX-CHOP, EVAP or EPOCH.

54. The composition of Claim 44, wherein the combination of
15 chemotherapeutic agents for treating NHL is: CVP, CHOP, C-MOPP, CAP-BOP, m-BACOD, ProMACE-MOPP, ProMACE-CytaBOM, MACOP-B, IMVP-16, MIME, DHAP, ESHAP, CEPP(B), or CAMP.

55. The composition of Claim 43, wherein the chemotherapeutic
20 agent for treating a B-cell leukemia is: anthracycline, cyclophosphamide, L-asparaginase, a purine analog.

56. The composition of Claim 43, wherein the combination of
25 chemotherapeutic agents for treating a B-cell leukemia is: vincristine, prednisone, anthracycline and cyclophosphamide or asparaginase; vincristine, prednisone, anthracycline, cyclophosphamide and asparaginase; CHOP; CMP; CVP; COP or CAP.

57. A method of treating a B cell malignancy in a subject in need of
30 such treatment comprising administering a therapeutically effective amount of at least one immunoregulating or immunomodulating antibody that is selected from the group consisting of an anti-CD23, anti-B7, anti-CD40, anti-CD40L and anti-CD4 antibody and at least B cell depleting antibody, and wherein said antibody administration is effected separately, in combination, and in either
35 order of administration.

5 58. The method of Claim 57 wherein the B cell depleting antibody is selected from the group consisting of an anti-CD19, anti-CD20, anti-CD22 and anti-CD37 antibody.

 59. The method of Claim 57 wherein B cell malignancy is non-
10 Hodgkin's lymphoma.

 60. The method of Claim 59 wherein said the NHL is selected from the subtype group consisting of: small lymphocytic, follicular and predominantly small cleaved cell, follicular and mixed small cleaved and large
15 cell type, follicular and predominantly large cell type, diffuse small cleaved cell, diffuse mixed small and large cell, diffuse large cell, large cell immunoblastic, lymphoblastic, small non-cleaved Burkitt's and non-Burkitt's type, AIDS-related lymphomas, angioimmunoblastic lymphadenopathy, mantle cell lymphoma, and monocytoid B-cell lymphoma.

20

 61. The method of Claim 60 wherein said NHL is high grade, low grade or intermediate grade.

 62. The method of Claim 60 wherein said B cell depleting antibody
25 is an anti-CD20 or anti-CD22 antibody.

 63. The method of Claim 62 wherein said anti-CD20 antibody is RITUXAN®.

30 64. The method of Claim 62 wherein said anti-CD20 antibody is a human or humanized antibody.

 65. The method of Claim 57 wherein the B cell malignancy is B cell lymphoma.

35

5 66. The method of Claim 1 wherein the B cell malignancy is a leukemia.

 67. The method of Claim 66 wherein said leukemia is chronic lymphocytic leukemia, acute lymphoblastic leukemia or chronic B cell
10 leukemia.

 68. The method of Claim 57 wherein treatment comprises the administration of an anti-B7 antibody and an anti-CD20 antibody.

15 69. The method of Claim 68 wherein the anti-CD20 is RITUXAN®.

 70. The method of Claim 68 wherein the anti-B7 antibody is a Primatized® antibody.

20 71. The method of Claim 70 wherein the anti-B7 antibody induces apoptosis of cancer cells.

 72. The method of Claim 57 wherein the immunoregulatory antibody is administered after the B cell depleting antibody.

25

 73. The method of Claim 57 wherein the immunoregulatory antibody is administered before the B cell depleting antibody.

 74. The method of Claim 57 wherein the B cell depleting antibody
30 and the immunoregulatory antibody are administered within about a month of each other.

 75. The method of Claim 57 wherein the B cell depleting antibody and the immunoregulatory antibody are administered within about one week of
35 each other.

5 76. The method of Claim 57 wherein the B cell depleting antibody and the immunoregulatory antibody are administered within about 1 day of each other.

 77. The method of Claim 57 wherein is used to treat a B cell
10 malignancy selected from the group consisting of relapsed Hodgkin's disease, resistant Hodgkin's disease high grade, low grade and intermediate grade non-Hodgkin's lymphomas, small lymphocytic/B cell chronic lymphocytic leukemia (SLL/B-CLL), lymphoplasmacytoid lymphoma (LPL), mantle cell lymphoma (MCL), follicular lymphoma (FL), diffuse large cell lymphoma (DLCL),
15 Burkitt's lymphoma (BL), AIDS-related lymphomas, monocytic B cell lymphoma, angioimmunoblastic lymphadenopathy, small lymphocytic; follicular, diffuse large cell; diffuse small cleaved cell; large cell immunoblastic lymphoblastoma; small, non-cleaved; Burkitt's and non-Burkitt's; follicular, predominantly large cell; follicular, predominantly small cleaved cell; and
20 follicular, mixed small cleaved and large cell lymphomas.

 78. The method of Claim 77 wherein said B cell malignancy is Hodgkin's disease.

25 79. The method of Claim 57 wherein either or both antibody is attached to a radiolabel.

 80. The method of Claim 57 which further comprises chemotherapy or radiation therapy.

30

 81. The method of Claim 57 which includes administration of a non-antibody antagonist specific to CD40L or B7.

 82. A method of treating non-Hodgkin's lymphoma comprising
35 administering separately or in combination a therapeutically effective amount of an antibody to B7 and a B cell depleting anti-CD20 or anti-CD22 antibody.

5 83. The method of Claim 82 wherein said anti-CD20 antibody is
RITUXAN®.

 84. The method of Claim 82 wherein said anti-B7 antibody does not
inhibit the interaction of B7 antigen with CTLA4.

10

 85. The method of Claim 84 wherein said antibody to B7 is a human,
humanized, primatized or primate antibody.

 86. The method of Claim 82 wherein said NHL is high grade, low
15 grade or intermediate grade.

 87. The method of Claim 82 which includes administration of a
radiolabeled antibody.

20 88. A method of treating leukemia comprising administering a
therapeutically effective amount of an anti-B7 antibody and a B cell depleting
antibody specific to CD20 or CD22.

 89. The method of Claim 88 wherein said anti-CD20 antibody is
25 RITUXAN® (antibody provided by ATCC 69119).

 90. The method of Claim 88 wherein said leukemia is chronic
lymphocytic leukemia, acute lymphoblastic leukemia or chronic B cell
leukemia.

30

 91. The method of Claim 88 wherein either or both of said
antibodies are chimeric, bispecific, human or humanized antibodies.

 92. The method of Claim 57 wherein the anti-B7 antibody is a
35 depleting antibody.

5 93. The method of Claim 57 wherein the anti-B7 antibody is a non-depleting antibody.

 94. The method of Claim 57 wherein the anti-B7 antibody specifically binds B7.1 (CD80).

10

 95. The method of Claim 57 wherein the anti-B7 antibody specifically binds B7.2 (CD86).

 96. The method of Claim 57 which includes administration of a
15 radiolabeled anti-CD22 or anti-CD22 antibody.

 97. The method of Claim 96 wherein said radiolabel is yttrium.

 98. The method of Claim 97 wherein said radiolabeled anti-CD20 is
20 yttrium-labeled RITUXAN® or yttrium-labeled 2B8.


 99. The method of Claim 82 wherein the anti-B7 antibody is a non-depleting antibody.

25 100. The method of Claim 82 wherein the anti-B7 antibody is a depleting antibody.

 101. The method of Claim 82 wherein the non-Hodgkin's lymphoma is selected from small lymphocytic, follicular and predominantly small cleaved
30 cell, follicular and mixed small cleaved and large cell type, follicular and predominantly large cell type, diffuse small cleaved cell, diffuse mixed small and large cell, diffuse large cell, large cell immunoblastic, lymphoblastic, small non-cleaved Burkitt's and non-Burkitt's type, AIDS-related lymphomas, angioimmunoblastic lymphadenopathy, mantle cell lymphoma, and monocytoid
35 B-cell lymphoma.


- 5 102. The method of Claim 57 which includes chemotherapy.
103. The method of Claim 82 which includes chemotherapy.

1/6

**FIG. 1**

2/6

ANTI-CD40L ANTIBODY (IDEC-131) OVERRIDES CD40L MEDIATED
RESISTANCE TO KILLING BY ADM OF B-LYMPHOMA CELLS



idéc-131

FIG. 2a

3/6

EFFECT OF RITUXAN ON NORMAL AND sCD40L PRE-TREATED DHL-4 CELLS




FIG. 2b

4/6

BLOCKING OF CD40L MEDIATED CELL SURVIVAL OF B-CLL BY ANTI-CD40L ANTIBODY (IDEC-131)




FIG. 3a

5/6




FIG. 3b

6/6




FIG. 4

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/15677

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 39/395; C07K 16/32

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST, DIALOG, BIOSIS, CA, EMBASE, MEDLINE
search terms: leukemia, lymphoma, cd40L, cd40 ligand, gp39, cd20, cd19, cd22, antibod?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,736,137 A (ANDERSON et al.) 07 April 1998, see entire document.	1-103
Y	US 6,001,358 A (BLACK et al.) 14 December 1999, see entire document.	1-103

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	* T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &* document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means	
* P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 JULY 2001

Date of mailing of the international search report

23 OCT 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PHILLIP GAMBEL

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1998) *

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/15677

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/1.49, 130.1, 133.1, 141.1, 143.1, 144.1, 153.1, 154.1, 173.1, 178.1 181.1, 183.1; 530/387.1, 387.3, 388.1, 388.2, 388.22, 388.75, 388.7, 388.73, 388.75, 391.1, 391.3, 391.7

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

424/1.49, 130.1, 133.1, 141.1, 143.1, 144.1, 153.1, 154.1, 173.1, 178.1 181.1, 183.1; 530/387.1, 387.3, 388.1, 388.2, 388.22, 388.75, 388.7, 388.73, 388.75, 391.1, 391.3, 391.7

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)